

*MATRIX FACTORIZATIONS FOR
DOMESTIC TRIANGLE SINGULARITIES*

BY

DAWID EDMUND KĘDZIERSKI (Szczecin), HELMUT LENZING (Paderborn)
and HAGEN MELTZER (Szczecin)

Abstract. Working over an algebraically closed field k of any characteristic, we determine the matrix factorizations for the—suitably graded—triangle singularities $f = x^a + y^b + z^c$ of domestic type, that is, we assume that (a, b, c) are integers at least two satisfying $1/a + 1/b + 1/c > 1$. Using work by Kussin–Lenzing–Meltzer, this is achieved by determining projective covers in the Frobenius category of vector bundles on the weighted projective line of weight type (a, b, c) . Equivalently, in a representation-theoretic context, we can work in the mesh category of $\mathbb{Z}\tilde{\Delta}$ over k , where $\tilde{\Delta}$ is the extended Dynkin diagram corresponding to the Dynkin diagram $\Delta = [a, b, c]$. Our work is related to, but in methods and results different from, the determination of matrix factorizations for the \mathbb{Z} -graded simple singularities by Kajiwara–Saito–Takahashi. In particular, we obtain symmetric matrix factorizations whose entries are scalar multiples of monomials, with scalars taken from $\{0, \pm 1\}$.

1. Introduction. Assuming (a, b, c) is a triple of integers greater than or equal to 2, we investigate the \mathbb{L} -graded hypersurface $S = k[x_1, x_2, x_3]/(f)$ determined by the triangle singularity $f = x_1^a + x_2^b + x_3^c$. Here, $\mathbb{L} = \mathbb{L}(a, b, c)$ is the rank one abelian group on generators $\vec{x}_1, \vec{x}_2, \vec{x}_3$ with relations $a\vec{x}_1 = b\vec{x}_2 = c\vec{x}_3 =: \vec{c}$, and the generator x_i from S is given degree \vec{x}_i ($i = 1, 2, 3$). Note that the polynomial f is homogeneous of degree \vec{c} , the *canonical element* of \mathbb{L} . Let $\mathbb{X} = \mathbb{X}(a, b, c)$ be the associated weighted projective line, whose category of coherent sheaves $\text{coh}(\mathbb{X})$ is obtained from S by Serre’s construction as the quotient category $\text{mod}^{\mathbb{L}}(S)/\text{mod}_0^{\mathbb{L}}(S)$ (see [GL87, Section 1.8]). Sheafification, given by the natural quotient functor $q : \text{mod}^{\mathbb{L}}(S) \rightarrow \text{coh}(\mathbb{X})$, then induces an equivalence between the full subcategory $\text{CM}^{\mathbb{L}}(S)$ of \mathbb{L} -graded (maximal) Cohen–Macaulay modules over S and the category $\text{vect}(\mathbb{X})$ of vector bundles on \mathbb{X} [GL87, Theorem 5.1]. Since S is graded Gorenstein, $\text{CM}^{\mathbb{L}}(S)$ is a Frobenius category with respect to the exact structure inherited from the abelian category $\text{mod}^{\mathbb{L}}(S)$ of finitely generated \mathbb{L} -graded

2010 *Mathematics Subject Classification*: Primary 14J17, 13H10, 16G60; Secondary 16G70.
Key words and phrases: triangle singularity, matrix factorization, weighted projective line, vector bundle, singularity category, Cohen–Macaulay module, projective cover, injective hull.

S -modules. With respect to this structure, the graded maximal Cohen–Macaulay modules of rank one form the indecomposable projective-injectives of $\mathrm{CM}^{\mathbb{L}}(S)$. The corresponding stable category $\underline{\mathrm{CM}}^{\mathbb{L}}(S)$ is triangulated. It is equivalent to the *singularity category* $\mathrm{Sing}^{\mathbb{L}}(S)$ introduced and studied by Buchweitz [Buc86] in the ungraded case and by Orlov [Orl09] in the graded case.

Important for the present paper is an alternative description of a singularity category as the *stable category of vector bundles* $\underline{\mathrm{vect}}(\mathbb{X})$ on the weighted projective line \mathbb{X} (see [KLM13]). To define this category, we call a sequence $\eta : 0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$ of vector bundles *distinguished-exact* if $\mathrm{Hom}(L, \eta)$ is exact for each line bundle L on \mathbb{X} . With the exact structure defined by these sequences, the category $\mathrm{vect}(\mathbb{X})$ of vector bundles on \mathbb{X} is a Frobenius category, equivalent to $\mathrm{CM}^{\mathbb{L}}(S)$, such that the indecomposable projective-injectives are just the line bundles on \mathbb{X} . A fortiori, the stable category $\underline{\mathrm{CM}}^{\mathbb{L}}(S)$ of Cohen–Macaulay modules is equivalent to the factor category $\underline{\mathrm{vect}}(\mathbb{X}) = \mathrm{vect}(\mathbb{X})/[\mathcal{L}]$, where $[\mathcal{L}]$ is the ideal consisting of all morphisms factoring through a finite direct sum of line bundles.

By results of Buchweitz [Buc86] and Orlov [Orl09], it is known that the singularity category $\mathrm{Sing}^{\mathbb{L}}(S)$, in the \mathbb{L} -graded sense, and the category of \mathbb{L} -graded maximal Cohen–Macaulay modules $\underline{\mathrm{CM}}^{\mathbb{L}}(S)$ are equivalent. Thus the stable category of vector bundles $\underline{\mathrm{vect}}(\mathbb{X})$ is another incarnation of the singularity category. In addition, all these categories are triangle equivalent to $\underline{\mathrm{MF}}^{\mathbb{L}}(T, f)$, the *stable category of \mathbb{L} -graded matrix factorizations* of f over the polynomial algebra $T = k[x_1, x_2, x_3]$.

For a base field of characteristic zero, a related category of graded matrix factorizations of a \mathbb{Z} -graded simple singularity was investigated by H. Kajiwara, K. Saito and A. Takahashi [KST07]. While these authors work directly inside the category of matrix factorizations, we work inside the category of vector bundles on the associated weighted projective line, and exploit well-known results on the Auslander–Reiten theory of $\mathrm{vect}(\mathbb{X})$. By contrast, our paper takes as a starting point the study of triangle singularities, and the associated stable category of vector bundles [KLM13]. Accordingly, we work over an algebraically closed field k of arbitrary characteristic. We recall that $\chi_{\mathbb{X}} = 1 - (1/a + 1/b + 1/c)$ is the *Euler characteristic* of \mathbb{X} such that domestic type for \mathbb{X} relates to positive Euler characteristic.

For a weighted projective line \mathbb{X} of domestic weight type (a, b, c) , the main achievement of our paper is two-fold: (A) a complete description of the projective covers (resp. the injective hulls) of indecomposable vector bundles, and (B) a complete description of all \mathbb{L} -graded matrix factorizations for singularities $f = x_1^a + x_2^b + x_3^c$ for indecomposable \mathbb{L} -graded (maximal) Cohen–Macaulay modules.

To achieve (A), we start with a fundamental result from [KLM13] on the projective covers, and likewise the injective hulls, of indecomposable vector bundles of rank two. For this first step there is no restriction on the Euler characteristic. Then in the second step, assuming domestic type, we use the knowledge of the Auslander–Reiten quiver for the category $\text{vect}(\mathbb{X})$, and use properly chosen distinguished-exact sequences to “extend” the projective covers to indecomposable bundles of higher rank. To achieve (B), we then lift minimal projective resolutions in $\text{CM}^{\mathbb{L}}(S) = \text{vect}(\mathbb{X})$ to matrix factorizations. As a key ingredient of the proof, we use the fact that the indecomposable vector bundles involved are uniquely determined by their projective covers (see Proposition 3.11).

We remark that step (A) has a direct interpretation in the representation theory of path algebras of extended Dynkin quivers: Assuming domestic type, it follows from a combination of [GL87] and [Hap88] that the category of indecomposable vector bundles on \mathbb{X} is equivalent to the mesh category $k(\mathbb{Z}\Delta)$ for the extended Dynkin star $\Delta = [a, b, c]$. Our results on projective covers and matrix factorizations thus offer new insight in the nature of the representation theory for path algebras of extended Dynkin type.

2. Basic concepts. We briefly recall the concept of a weighted projective line, where we restrict to the case of *triple weight type*, given by weight triples (a, b, c) of integers greater than or equal to 2. For a more general setting and further details we refer to [GL87]. Throughout the present paper, k denotes an algebraically closed field.

Let $\mathbb{L} = \mathbb{L}(a, b, c)$ be the rank one abelian group on generators $\vec{x}_1, \vec{x}_2, \vec{x}_3$ with relations $a\vec{x}_1 = b\vec{x}_2 = c\vec{x}_3 =: \vec{c}$, where \vec{c} is called the *canonical element*. We note that \mathbb{L} is naturally isomorphic to the *Picard group* of \mathbb{X} . The polynomial algebra $T = k[x_1, x_2, x_3]$, $T = T(a, b, c)$, is equipped with an \mathbb{L} -grading by giving x_i degree \vec{x}_i for each $i = 1, 2, 3$. Further, let $S = S(a, b, c)$ denote the factor algebra $k[x_1, x_2, x_3]/(f)$, where $f = x_1^a + x_2^b + x_3^c$. Because f is a homogeneous polynomial, the algebra S is also \mathbb{L} -graded; by $S_{\vec{x}}$ we denote the finite-dimensional vector space of elements of degree \vec{x} . The weighted projective line $\mathbb{X} = \mathbb{X}(a, b, c)$ is by definition the \mathbb{L} -graded projective spectrum of the \mathbb{L} -graded algebra S . By [GL87] its category of coherent sheaves $\text{coh}(\mathbb{X})$ is obtained by Serre’s construction as the quotient category of $\text{mod}^{\mathbb{L}}(S)$, the category of finitely generated \mathbb{L} -graded S -modules, by the Serre subcategory $\text{mod}_0^{\mathbb{L}}(S)$ of all finite-dimensional (= finite length) modules. By $q : \text{mod}^{\mathbb{L}}(S) \rightarrow \text{coh}(\mathbb{X})$, $M \mapsto \widetilde{M}$, we denote the corresponding quotient functor (sheafification).

For the present paper, the following result is of importance. For its proof, we refer to [GL87, Theorem 5.1], and for the last claim to [KLM13]. From the

first paper we take the following description of \mathbb{L} -graded (maximal) Cohen–Macaulay modules: A finitely generated \mathbb{L} -graded S -module M is (maximal) Cohen–Macaulay if and only if $\text{Hom}(k(\vec{x}), M) = 0 = \text{Ext}^1(k(\vec{x}), M)$ for each $\vec{x} \in \mathbb{L}$. Hereafter, Cohen–Macaulay will always mean *maximal* \mathbb{L} -graded Cohen–Macaulay.

PROPOSITION 2.1. *The sheafification functor $q : \text{mod}^{\mathbb{L}}(S) \rightarrow \text{coh}(\mathbb{X})$, $M \mapsto \tilde{M}$, induces an equivalence $q : \text{CM}^{\mathbb{L}}(S) \xrightarrow{\sim} \text{vect}(\mathbb{X})$ between the category $\text{CM}^{\mathbb{L}}(S)$ of finitely generated \mathbb{L} -graded S -modules and the category $\text{vect}(\mathbb{X})$ of vector bundles on \mathbb{X} . This functor also induces an equivalence between the full subcategories $\text{proj}^{\mathbb{L}}(S)$ of finitely generated \mathbb{L} -graded projective modules and the category \mathcal{L} of line bundles on \mathbb{X} . Accordingly, q induces a triangle equivalence between the stable categories $\underline{\text{CM}}^{\mathbb{L}}(S)$ and $\underline{\text{vect}}(\mathbb{X})$. ■*

We now collect some facts on $\text{coh}(\mathbb{X})$. This category is hereditary, that is, all extensions of degree ≥ 2 vanish, and it admits Serre duality in the form $\text{DExt}_{\mathbb{X}}^1(F, G) \simeq \text{Hom}_{\mathbb{X}}(G, \tau_{\mathbb{X}}F)$, where D denotes the usual duality $\text{Hom}_k(-, k)$ and $\tau_{\mathbb{X}}F = F(\vec{\omega})$, where $\vec{\omega} = \vec{c} - \sum_{i=1}^3 \vec{x}_i$ is the *dualizing element* of \mathbb{L} . Consequently, $\text{coh}(\mathbb{X})$ has almost-split sequences, and the Auslander–Reiten translation $\tau_{\mathbb{X}}$ is a self-equivalence of $\text{coh}(\mathbb{X})$ given by the degree shift $X \mapsto X(\vec{\omega})$.

The complexity of the classification problem of vector bundles on \mathbb{X} is largely determined by the *Euler characteristic* of \mathbb{X} , given by the expression $\chi_{\mathbb{X}} = 1/a + 1/b + 1/c - 1$. A weighted projective line \mathbb{X} is said to be of *domestic type* if $\chi_{\mathbb{X}} > 0$. Consequently, in our setup, $\mathbb{X}(a, b, c)$ is of domestic type if and only if the weight type is, up to permutation, one of the following: $(2, 2, n)$ ($n \geq 2$), $(2, 3, 3)$, $(2, 3, 4)$, or $(2, 3, 5)$.

The concept of matrix factorizations was introduced by D. Eisenbud [Eis80]. For a textbook treatment we refer to [Yos90]. We recall the definition and some basic facts, adapted to the present \mathbb{L} -graded setting. Let $T = k[x_1, x_2, x_3]$ be the polynomial algebra, viewed as \mathbb{L} -graded algebra, and fix the polynomial $f = x_1^a + x_2^b + x_3^c$. An *\mathbb{L} -graded matrix factorization of f* is a pair of homogeneous T -linear maps $\varphi : P_1 \rightarrow P_0$ and $\psi : P_0 \rightarrow P_1(\vec{c})$ for \mathbb{L} -graded projective T -modules P_0 and P_1 , denoted

$$(2.1) \quad P_1 \begin{array}{c} \xrightarrow{\varphi} \\ \xleftarrow{\psi} \end{array} P_0,$$

such that the compositions $\varphi\psi(-\vec{c}) : P_0(-\vec{c}) \rightarrow P_0$ and $\psi\varphi : P_1 \rightarrow P_1(\vec{c})$ are both the multiplication maps with f . Since P_0 and P_1 are \mathbb{L} -graded free T -modules, we may think of φ and ψ and $f\mathbf{1}$ as matrices whose entries are homogeneous members of T such that the two factorization conditions translate to the matrix equation $\varphi\psi = f\mathbf{1} = \psi\varphi$. We note that the degree shifts involved will mostly be clear from the context. For the matrix description

of a matrix factorization (2.1), we always assume that the decompositions of P_0 and $P_0(-\vec{c})$ (respectively P_1 and $P_1(\vec{c})$) into projectives of rank one are *compatible*, that is, correspond to each other by degree shift $X \mapsto X(-\vec{c})$. We note further that when describing a matrix factorization by matrices, we need to keep track of the direct sum decompositions of P_0 and P_1 into line bundles.

For two matrix factorizations $P_1 \begin{smallmatrix} \varphi \\ \leftarrow \\ \psi \end{smallmatrix} P_0$ and $P'_1 \begin{smallmatrix} \varphi' \\ \leftarrow \\ \psi' \end{smallmatrix} P'_0$, a pair (F_1, F_0) of (homogeneous) T -linear maps is called a *morphism of matrix factorizations* if the diagram

$$\begin{array}{ccccc} P_0(-\vec{c}) & \xrightarrow{\psi(-\vec{c})} & P_1 & \xrightarrow{\varphi} & P_0 \\ \downarrow F_0(-\vec{c}) & & \downarrow F_1 & & \downarrow F_0 \\ P'_0(-\vec{c}) & \xrightarrow{\psi'(-\vec{c})} & P'_1 & \xrightarrow{\varphi'} & P'_0 \end{array}$$

is commutative. Thinking of F_1 and F_0 (and also φ and ψ) as matrices whose entries are *homogeneous elements from T* , the two commutativity conditions (2.1) translate to matrix equations $F_0\varphi = \varphi'F_1$ and $F_1\psi = \psi'F_0$. We remark that a matrix factorization $P_1 \begin{smallmatrix} \varphi \\ \leftarrow \\ \psi \end{smallmatrix} P_0$ is indecomposable if and only if its endomorphism ring is local.

For any \mathbb{L} -graded matrix factorization (2.1), the cokernel $M = \text{cok}(P_1 \xrightarrow{\varphi} P_0)$ is annihilated by f , hence belongs to $\text{mod}^{\mathbb{L}}(S)$. Actually, M belongs to $\text{CM}^{\mathbb{L}}(S)$, and is called the (maximal) graded *Cohen–Macaulay S -module determined by (φ, ψ)* , also denoted as $\text{cok}(\varphi, \psi)$. Let $\text{MF}^{\mathbb{L}}(T, f)$ denote the category of all \mathbb{L} -graded matrix factorizations of f over T . Let \mathcal{U} denote the full subcategory of trivial matrix factorizations $(1, f)$. Then the assignment $(\varphi, \psi) \mapsto \text{cok}(\varphi)$ establishes an equivalence between the factor category $\text{MF}^{\mathbb{L}}(T, f)/[\mathcal{U}]$ and the category $\text{CM}^{\mathbb{L}}(S)$ of \mathbb{L} -graded Cohen–Macaulay modules over S . By means of the equivalence $q : \text{CM}^{\mathbb{L}}(S) \rightarrow \text{vect}(\mathbb{X})$, we may as well speak of the *vector bundle $E = q(\text{cok}(\varphi))$ determined by the matrix factorization (φ, ψ)* . For any projective T -module, the functor $\text{cok} : \text{MF}^{\mathbb{L}}(T, f) \rightarrow \text{CM}^{\mathbb{L}}(S)$ sends $P \begin{smallmatrix} \xrightarrow{1} \\ \leftarrow \\ f \end{smallmatrix} P$ to zero and $P(-\vec{c}) \begin{smallmatrix} \xrightarrow{f} \\ \leftarrow \\ 1 \end{smallmatrix} P$ to the projective S -module $\bar{P} = P/f.P$, and all projective S -modules are obtained in this way.

We are now in a position to formulate Eisenbud’s matrix factorization theorem [Eis80], adapted to our \mathbb{L} -graded context. We follow Yoshino’s presentation [Yos90].

THEOREM 2.2. *Let \mathcal{U} , respectively $\bar{\mathcal{U}}$, be the full subcategory of $\text{MF}^{\mathbb{L}}(T, f)$ consisting of all $P \begin{smallmatrix} \xrightarrow{1} \\ \leftarrow \\ f \end{smallmatrix} P$, respectively of all matrix factorizations $P \begin{smallmatrix} \xrightarrow{1} \\ \leftarrow \\ f \end{smallmatrix} P$*

and $P(-\vec{c}) \xrightarrow[1]{f} S$. Then the functor $\text{cok} : \text{MF}^{\mathbb{L}}(T, f) \rightarrow \text{CM}^{\mathbb{L}}(S)$ induces equivalences

$$\text{MF}^{\mathbb{L}}(T, f)/[\mathcal{U}] \rightarrow \text{CM}^{\mathbb{L}}(S) \quad \text{and} \quad \text{MF}^{\mathbb{L}}(T, f)/[\overline{\mathcal{U}}] \rightarrow \underline{\text{CM}}^{\mathbb{L}}(S).$$

Moreover, the suspension functor of the triangulated category $\underline{\text{CM}}^{\mathbb{L}}(S)$ is induced by the (functorial) expression $(P_1 \xrightleftharpoons[\psi]{\varphi} P_0)[1] = P_0 \xrightleftharpoons[\varphi]{\psi} P_1(\vec{c})$. ■

We say that a matrix factorization $P_0 \xrightleftharpoons[\psi]{\varphi} P_1$ is *reduced* if φ and ψ belong to the radical of $\text{mod}^{\mathbb{L}}(T)$, that is, if viewed as matrices, φ and ψ have entries in the graded maximal ideal (x_1, x_2, x_3) of T . The cokernel M of a reduced matrix factorization is an \mathbb{L} -graded Cohen–Macaulay module over $S = T/(f)$ without projective summands; moreover, iterating the formation of matrix factorizations of f over T , we obtain a sequence

$$(2.2) \quad \cdots \xrightarrow{\psi} P_1(-\vec{c}) \xrightarrow{\varphi} P_0(-\vec{c}) \xrightarrow{\psi} P_1 \xrightarrow{\varphi} P_0 \rightarrow M \rightarrow 0$$

of matrix factorizations which is 2-periodic up to degree shift with \vec{c} . Reduction modulo (f) then yields the sequence

$$(2.3) \quad \cdots \xrightarrow{\bar{\psi}} \bar{P}_1(-\vec{c}) \xrightarrow{\bar{\varphi}} \bar{P}_0(-\vec{c}) \xrightarrow{\bar{\psi}} \bar{P}_1 \xrightarrow{\bar{\varphi}} \bar{P}_0 \rightarrow M \rightarrow 0,$$

which is a minimal \mathbb{L} -graded projective and 2-periodic resolution of M over S . Here, the bar always stands for reduction modulo f .

In order to determine a matrix factorization $P_1 \xrightleftharpoons[\psi]{\varphi} P_0$ for a Cohen–Macaulay module M without projective summands, we will first determine the minimal projective resolution (2.3) of M over S , and then lift the S -matrix pair $(\bar{\varphi}, \bar{\psi})$ to a matrix pair (φ, ψ) over T , such that *additionally* $\varphi\psi = f\mathbb{1} = \psi\varphi$.

For weight triples $(2, a, b)$, the suspension functor [1] for $\underline{\text{vect}}(\mathbb{X})$ is induced by the degree shift $X \mapsto X(\vec{x}_1)$ by \vec{x}_1 (see [KLM13, Proposition 6.8]). This allows us to introduce the concept of a *symmetric matrix factorization* of f for a graded Cohen–Macaulay module M without projective summands (correspondingly for a vector bundle E without line bundle summands).

Namely, we call a matrix factorization $P_1 \xrightleftharpoons[\psi]{\varphi} P_0$ for M *symmetric*—recall that then $\varphi : P_1 \rightarrow P_0$ and $\psi : P_0 \rightarrow P_1(\vec{c})$ are homogeneous T -linear maps—provided $P_0 = P_1(\vec{x}_1)$ and $\psi = \varphi(\vec{x}_1)$. Note that this requirement makes sense since $2\vec{x}_1 = \vec{c}$. In this case, by abuse of notation, we will—as for ungraded symmetric matrix factorizations—also write $\varphi = \psi$. We will show in Section 5 that for weight type $(2, a, b)$, each indecomposable vector bundle of rank two is determined by a symmetric matrix factorization of f .

Moreover, if we deal with domestic type, necessarily given by a weight triple $(2, a, b)$, then each indecomposable vector bundle of rank at least two will admit a symmetric matrix factorization by Theorem 5.4.

For a weight triple (p_1, p_2, p_3) we set $\bar{p} = \text{l.c.m.}(p_1, p_2, p_3)$. There is a unique group homomorphism $\delta : \mathbb{L} \rightarrow \mathbb{Z}$ called the *degree map* which sends \vec{x}_i to \bar{p}/p_i . The kernel of δ is the torsion group $t\mathbb{L}$ of \mathbb{L} . Further, there is a unique group homomorphism $\text{deg} : K_0(\text{coh}(\mathbb{X})) \rightarrow \mathbb{Z}$, called the *degree*, such that $\text{deg}([\mathcal{O}(\vec{x})]) = \delta(\vec{x})$ for each $\vec{x} \in \mathbb{L}$. For each non-zero $X \in \text{coh}(\mathbb{X})$ at least one of $\text{rk}(X)$ or $\text{deg}(X)$ is non-zero, yielding a well defined *slope* $\mu(X) = \text{deg}(X)/\text{rk}(X)$ in the extended rationals $\mathbb{Q} \cup \{\infty\}$. The slope of an indecomposable object X is a useful indicator of the position of X in the category $\text{coh}(\mathbb{X})$. In the domestic situation, moreover, each indecomposable vector bundle X is *stable*, that is, satisfies $\mu(X') < \mu(X)$ for each proper subobject $0 \neq X' \subsetneq X$. Still assuming domestic type, stability of a non-zero vector bundle X implies $\text{End}(X) = k$ and $\text{Ext}_{\mathbb{X}}^1(X, X) = 0$, that is, the exceptionality of X . For all foregoing facts see [GL87].

Sometimes a refinement of the degree, called *determinant*, is necessary. This is a group homomorphism $\det : K_0(\text{coh}(\mathbb{X})) \rightarrow \mathbb{L}$ such that $\det(\mathcal{O}(\vec{x})) = \vec{x}$ for each $\vec{x} \in \mathbb{L}$. In particular, $\text{deg} = \delta \circ \det$ (see [LM92, 2.7]).

By means of a line bundle filtration for a vector bundle E one further obtains the formula


$$(2.4) \quad \det(E(\vec{x})) = \det(E) + \text{rk}(E) \cdot \vec{x} \quad \text{for all } \vec{x} \in \mathbb{L}.$$

We finally recall from [GL87] that the category $\text{coh}(\mathbb{X})$ has almost split sequences with the Auslander–Reiten translation given by degree shift $X \mapsto X(\vec{\omega})$ with the dualizing element $\vec{\omega} = \vec{c} - \sum_{i=1}^3 \vec{x}_i$.

The category of vector bundles for domestic weight triples. Recall that a weight triple (a, b, c) with entries ≥ 2 has domestic type if and only if it is one of $(2, 2, n)$, $n \geq 2$, $(2, 3, 3)$, $(2, 3, 4)$, or $(2, 3, 5)$. The shape of the Auslander–Reiten quiver of $\text{vect}(\mathbb{X})$ is then $\mathbb{Z}\Delta$, where Δ is the extended Dynkin diagram, attached to the Dynkin star $[a, b, c]$. The category of indecomposable vector bundles is then equivalent to the mesh category $k(\mathbb{Z}\Delta)$. In this case, the stable category $\underline{\text{vect}}(\mathbb{X})$ is equivalent to the bounded derived category $D^b(\text{mod}(A))$ for the path algebra $A = kQ$ of Dynkin type Δ' obtained from Δ by removing all vertices where the standard additive function on Δ takes value 1 [KLM13, Section 5.1]. The table below summarizes the situation.

Weight triple	$(2, 2, n)$	$(2, 3, 3)$	$(2, 3, 4)$	$(2, 3, 5)$
Δ	$\tilde{\mathbb{D}}_{n+2}$	$\tilde{\mathbb{E}}_6 = [3, 3, 3]$	$\tilde{\mathbb{E}}_7 = [2, 4, 4]$	$\tilde{\mathbb{E}}_8 = [2, 3, 6]$
Δ'	$\mathbb{A}_{n-1} = [n-1]$	$\mathbb{D}_4 = [2, 2, 2]$	$\mathbb{E}_6 = [2, 3, 3]$	$\mathbb{E}_8 = [2, 3, 5]$

For the rest of the paper, it is important to understand how the Picard group \mathbb{L} acts on the mesh category $k(\mathbb{Z}\Delta)$, or on the underlying translation quiver $\mathbb{Z}\Delta$, by degree shift. We illustrate this for the weight triple $(2, 3, 4)$, where a piece of the Auslander–Reiten quiver is depicted below. The considerations are similar for other domestic weight triples. We first remark that the rank of vector bundles is constant on τ -orbits; the values of the rank are displayed at the right end.



We thus have two τ -orbits of line bundles, the lower and the upper border, three τ -orbits of indecomposable rank-two bundles, two τ -orbits of indecomposable bundles of rank 3 and a single τ -orbit of rank 4. Since the Picard group acts transitively on the iso-classes of line bundles, we may freely choose the position of the structure sheaf from one of the two line bundle orbits. Once this is done, the position of the other line bundles is fixed, up to a symmetry of $\mathbb{Z}\Delta$. To indicate the position of a line bundle $\mathcal{O}(\vec{x})$, we use the bracket notation (\vec{x}) such that the structure sheaf is given by the symbol $(\vec{0})$, and its Auslander–Reiten translate $\tau\mathcal{O}$ is given by the symbol $(\vec{\omega})$, where $\vec{\omega} = \vec{c} - (\vec{x}_1 + \vec{x}_2 + \vec{x}_3)$ and hence $\delta(\vec{\omega}) = -1$. This now allows us to determine easily the values of the degree function for each indecomposable vector bundle. Since $\mathcal{O}(\vec{x}_3)$ has degree 3 and $\text{Hom}_{\mathbb{X}}(\mathcal{O}, \mathcal{O}(\vec{x}_3)) = k$, there is only one choice for the position (\vec{x}_3) , once the position $(\vec{0})$ has been fixed. All further line bundles are then given by one of the symbols $(\vec{0} + n\vec{\omega})$, respectively $(\vec{x}_3 + n\vec{\omega})$, with $n \in \mathbb{Z}$.

Corresponding to the positions (\vec{x}_1) , (\vec{x}_2) and (\vec{x}_3) in the mesh category, the shift actions by \vec{x}_1 , \vec{x}_2 and \vec{x}_3 are given as follows: The shift by \vec{x}_1 (resp. \vec{x}_3) is a glide reflection, composed with reflection with respect to the central horizontal axis with the sixth resp. third power of τ^- . Further, the shift action by \vec{x}_2 equals τ^{-4} .

Finally, let us remark that, obviously, the factor category $\text{vect}(\mathbb{X})/[\mathcal{L}]$ obtained from $\text{vect}(\mathbb{X})$ for $\mathbb{X} = \mathbb{X}(2, 3, 4)$ by factoring out the two line bundle orbits yields the mesh category $k(\mathbb{Z}\mathbb{E}_6)$, equivalent to $D^b(\text{mod}(kQ))$ for a quiver Q of type \mathbb{E}_6 , thus illustrating the facts mentioned at the beginning of this section.

REMARK 2.3. In view of Theorem 3.3, it is useful to interpret the degree shift by \vec{x}_1 in terms of the Auslander–Reiten quiver of $\text{vect}(\mathbb{X})$ resp. $\underline{\text{vect}}(\mathbb{X})$. For this, we assume domestic type $(2, a, b)$.

- For type $(2, 3, 5)$ we have $\vec{x}_1 = -15\vec{\omega}$. Thus the degree shift by \vec{x}_1 is translation to the right by 15 mesh units.

- For type $(2, 3, 4)$ we have $\vec{x}_1 = -6\vec{\omega} + (\vec{x}_1 - 2\vec{x}_3)$. We note that the element $\vec{x}_1 - 2\vec{x}_3$ has order two. Thus the degree shift by \vec{x}_1 is the glide reflection given by composing reflection in the central axis with translation to the right by 6 mesh units.

- For type $(2, 3, 3)$, we have $\vec{x}_1 = -3\vec{\omega}$, and the degree shift by \vec{x}_1 is translation to the right by 3 mesh units.

- For type $(2, 2, n)$, we use the fact that the degree shifts by \vec{x}_1 , \vec{x}_2 and $-\vec{\omega}$ agree on objects of $\underline{\text{vect}}(\mathbb{X})$ [KLM13, p. 235], and only deal with the shift action of \vec{x}_1 on objects of $\text{vect}(\mathbb{X})$. Further, we need to distinguish whether n is even or odd: For $n = 2k$ (resp. $n = 2k + 1$) the degree shifts by $-k\vec{\omega}$ and $\vec{x}_1 + (k\vec{x}_3 - \vec{x}_1)$ (resp. by $-k\vec{\omega}$ and $\vec{x}_1 + (k\vec{x}_3 - \vec{x}_1)$) agree on objects of $\underline{\text{vect}}(\mathbb{X})$. In the first case the element $k\vec{x}_3 - \vec{x}_1$ has order two, while in the second case we obtain $2(k\vec{x}_3 - \vec{x}_1) = -\vec{x}_3$. Hence the degree shift by \vec{x}_1 on the Auslander–Reiten quiver of $\underline{\text{vect}}(\mathbb{X})$ is the glide reflection given by composing reflection in the central axis with translation to the right by k mesh units (resp. by $k + 1/2$ mesh units).

3. Projective covers. When speaking of weight triples, we always assume that the weights are at least two. In the domestic case this just excludes the weight types $(\)$, (a) and (a, b) where each indecomposable vector bundle is a line bundle, and the matrix factorization problem thus becomes trivial.

General results. Assuming an arbitrary weight triple (p_1, p_2, p_3) , this section starts by quoting two general results [KLM13, Theorems 4.2 and 4.6] on indecomposable vector bundles of rank two and their projective covers in $\text{vect}(\mathbb{X})$. We recall that the double suspension functor for $\underline{\text{vect}}(\mathbb{X})$ is induced by degree shift with the canonical element \vec{c} . Moreover, for weight triples $(2, a, b)$, the suspension functor itself is induced by the degree shift with \vec{x}_1 [KLM13, Proposition 6.8]. Switching now to weight triples of domestic type, necessarily of type $(2, a, b)$, the aim of this section is to determine the projective cover (and likewise the injective hull) for each indecomposable vector bundle of rank ≥ 2 .

We assume triple weight type (p_1, p_2, p_3) . Let $\delta = \vec{c} + 2\vec{\omega}$ be the *dominant element* of \mathbb{L} . The elements $\vec{0} \leq \vec{x} \leq \vec{\delta}$ then have the form $\vec{x} = \sum_{i=1}^3 l_i \vec{x}_i$ with $0 \leq l_i \leq p_i - 2$. Following [KLM13, Section 4], a vector bundle E of

rank 2 is called an *extension bundle* if E is the middle term of a non-split exact sequence

$$(3.1) \quad \eta_{\vec{x}} : 0 \rightarrow L(\vec{\omega}) \rightarrow E \rightarrow L(\vec{x}) \rightarrow 0,$$

where L is a line bundle and $\vec{0} \leq \vec{x} \leq \vec{\delta}$. Because $\text{Ext}_{\mathbb{X}}^1(L(\vec{x}), L(\vec{\omega})) = k$, the bundle E is uniquely determined, up to isomorphism; we then denote E by $E_L\langle\vec{x}\rangle$. For $L = \mathcal{O}$ we just write $E\langle\vec{x}\rangle$. If $\vec{x} = 0$, then the sequence $\eta_{\vec{x}}$ is almost-split, and $E = E_L\langle 0 \rangle$ is called an *Auslander bundle*, more precisely the Auslander bundle attached to L . Applying degree shift by \vec{y} from \mathbb{L} to the exact sequence (3.1), we obtain the useful identity

$$(3.2) \quad (E_L\langle\vec{x}\rangle)(\vec{y}) \cong E_{L(\vec{y})}\langle\vec{x}\rangle \quad \text{for all } 0 \leq \vec{x} \leq \vec{\delta}, \vec{y} \in \mathbb{L}.$$

We recall that an object E in an abelian (resp. a triangulated) category is *exceptional* if $\text{End}(E) = k$ and further $\text{Ext}^d(E, E) = 0$ (resp. $\text{Hom}(E, E[d]) = 0$) for each integer $d \neq 0$. For objects of a hereditary category, like $\text{coh}(\mathbb{X})$, the Ext-condition only has to be checked for $d = 1$.

The following three theorems from [KLM13] mark the starting point of our investigation. For the first one we refer to Theorem 4.2 and Corollary 4.11 from the cited paper, and for the second one to Theorem 4.6 there. We recall that $\vec{\delta} = \sum_{i=1}^3 (p_i - 2)\vec{x}_i$ denotes the dominant element of \mathbb{L} .

THEOREM 3.1 (Vector bundles of rank two). *Assume \mathbb{X} is given by a weight triple (p_1, p_2, p_3) . Then:*

- (i) *Each indecomposable vector bundle of rank two is isomorphic to an extension bundle $E_L\langle\vec{x}\rangle$ for a suitable choice of a line bundle L and an element \vec{x} from \mathbb{L} satisfying $0 \leq \vec{x} \leq \vec{\delta}$.*
- (ii) *Each indecomposable vector bundle of rank two is exceptional in the category $\text{coh}(\mathbb{X})$ of coherent sheaves on \mathbb{X} . It is also exceptional in the stable category of vector bundles $\underline{\text{vect}}(\mathbb{X})$. ■*

THEOREM 3.2 (Projective and injective covers). *Assume \mathbb{X} is given by the weight triple (p_1, p_2, p_3) . Let $E_L\langle\vec{x}\rangle$, $0 \leq \vec{x} \leq \vec{\delta}$, be an extension bundle. Then its injective hull $\mathfrak{I}(E_L\langle\vec{x}\rangle)$ and its projective cover $\mathfrak{P}(E_L\langle\vec{x}\rangle)$ are given by*

$$(3.3) \quad \mathfrak{I}(E_L\langle\vec{x}\rangle) = L(\vec{x}) \oplus \bigoplus_{i=1}^3 L((1 + l_i)\vec{x}_i + \vec{\omega}),$$

$$(3.4) \quad \mathfrak{P}(E_L\langle\vec{x}\rangle) = L(\vec{\omega}) \oplus \bigoplus_{i=1}^3 L(\vec{x} - (1 + l_i)\vec{x}_i),$$

where $\vec{x} = l_1\vec{x}_1 + l_2\vec{x}_2 + l_3\vec{x}_3$.

Further, the four line bundle summands $(L_i)_{i=0}^3$ of $\mathfrak{I}(E_L\langle\vec{x}\rangle)$ (resp. $\mathfrak{P}(E_L\langle\vec{x}\rangle)$) are mutually Hom-orthogonal, that is,

$$\mathrm{Hom}(L_i, L_j) = \begin{cases} k & \text{if } i = j, \\ 0 & \text{if } i \neq j. \blacksquare \end{cases}$$

The next result is a straightforward consequence of [KLM13, Proposition 6.8].

THEOREM 3.3 (Weight type $(2, a, b)$). *Let \mathbb{X} be the weighted projective line of type $(2, a, b)$ and E be an indecomposable vector bundle of rank at least 2. There is a distinguished short exact sequence*

$$0 \rightarrow E(-\vec{x}_1) \rightarrow \mathfrak{P}(E) \xrightarrow{\pi_E} E \rightarrow 0,$$

where $\mathfrak{P}(E)$ is the projective cover of E , and likewise for the injective hull $\mathfrak{I}(E(-\vec{x}_1))$ of $E(-\vec{x}_1)$. In particular, $\mathfrak{I}(E) = \mathfrak{P}(E)(\vec{x}_1)$ and $\mathrm{rk}(\mathfrak{P}(E)) = 2\mathrm{rk}(E)$. \blacksquare

The following variant of the ‘horse-shoe lemma’ from homological algebra will be used to determine projective covers for vector bundles of larger rank. A dual result is valid for injective hulls.

LEMMA 3.4. *We assume weight type $(2, a, b)$. Let X and Y be vector bundles with projective covers $\mathfrak{P}(X) \xrightarrow{\pi_X} X$ and $\mathfrak{P}(Y) \xrightarrow{\pi_Y} Y$. Let*

$$(\star) \quad 0 \rightarrow X \xrightarrow{f} E \xrightarrow{g} Y \rightarrow 0$$

be a distinguished exact sequence in $\mathrm{vect}(\mathbb{X})$. (This condition is satisfied if (\star) is exact and $\mathrm{Ext}_{\mathbb{X}}^1(\mathfrak{P}(Y), X)$ is zero.) Then π_Y lifts to a map $\pi_Y^* : \mathfrak{P}(Y) \rightarrow E$ yielding a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathfrak{P}(X) & \longrightarrow & \mathfrak{P}(X) \oplus \mathfrak{P}(Y) & \longrightarrow & \mathfrak{P}(Y) & \longrightarrow & 0 \\ & & \pi_X \downarrow & & [f\pi_X, \pi_Y^*] \downarrow & & \pi_Y \downarrow & & \\ 0 & \longrightarrow & X & \xrightarrow{f} & E & \xrightarrow{g} & Y & \longrightarrow & 0 \end{array}$$

which establishes $\mathfrak{P}(X) \oplus \mathfrak{P}(Y)$ as a projective cover of E . Moreover, the rows of the diagram are distinguished exact and the vertical maps are distinguished epimorphisms.

Proof. We show that the condition $\mathrm{Ext}_{\mathbb{X}}^1(\mathfrak{P}(Y), X) = 0$ implies that (\star) is distinguished exact. Indeed, applying the functor $\mathrm{Hom}_{\mathbb{X}}(\mathfrak{P}(Y), -)$ to the exact sequence (\star) , we obtain a short exact sequence

$$\begin{aligned} 0 \rightarrow \mathrm{Hom}_{\mathbb{X}}(\mathfrak{P}(Y), X) &\xrightarrow{f^{\circ-}} \mathrm{Hom}_{\mathbb{X}}(\mathfrak{P}(Y), E) \xrightarrow{g^{\circ-}} \mathrm{Hom}_{\mathbb{X}}(\mathfrak{P}(Y), Y) \\ &\rightarrow \mathrm{Ext}_{\mathbb{X}}^1(\mathfrak{P}(Y), X) = 0 \end{aligned}$$

showing that each morphism from $\mathfrak{P}(Y)$ to X lifts to E . By definition of the projective cover, each morphism from a line bundle L to Y lifts to $\mathfrak{P}(Y)$.

Putting things together, any morphism from a line bundle L to Y lifts to a map $L \rightarrow E$, showing that (\star) is distinguished exact. Any of the two assumptions thus ensures that π_Y lifts to a map $\alpha : \mathfrak{P}(Y) \rightarrow E$ yielding the above diagram, and its claimed properties, by standard arguments of (relative) homological algebra.

It remains to explain why the minimality condition for the distinguished epimorphism $\pi_E : \mathfrak{P}(X) \oplus \mathfrak{P}(Y) \rightarrow E$ holds. Because of weight type $(2, a, b)$, we know from Theorem 3.3 that $\mathfrak{P}(E)$ and $\mathfrak{P}(X) \oplus \mathfrak{P}(Y)$ have the same rank $2 \operatorname{rk}(E)$, which ensures the claim. ■

We keep assuming weight type $(2, a, b)$. To obtain minimal projective resolutions, we have to determine those morphisms that are compositions of a projective cover $\mathfrak{P}(E) \xrightarrow{\pi_E} E$ with the corresponding injective hull $E \xrightarrow{j_E} \mathfrak{I}(E)$. The resulting morphism $u_E : \mathfrak{P}(E) \rightarrow \mathfrak{I}(E)$, $u_E = j_E \pi_E$, will be called a *cover morphism* for E . (Note that such cover morphisms depend on the projective cover and injective hull chosen.) Again, we reduce the determination of cover morphisms to the case of smaller rank.

LEMMA 3.5. *Let \mathbb{X} be of weight type $(2, a, b)$. Let (\star) be as in Lemma 3.4. Let \bar{u}_X (respectively \bar{u}_Y) be a cover morphisms for X (respectively Y). Then we obtain a cover morphism for E having the shape*

$$\bar{u}_E = \left[\begin{array}{c|c} \bar{u}_X & \beta \circ \alpha \\ \hline 0 & \bar{u}_Y \end{array} \right]$$

where $g \circ \alpha = \pi_Y$ and $\beta \circ f = j_X$.

Proof. From Lemma 3.4 we obtain the following commutative diagram with exact rows:


$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathfrak{P}(X) & \xrightarrow{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} & \mathfrak{P}(X) \oplus \mathfrak{P}(Y) & \xrightarrow{[0 \ 1]} & \mathfrak{P}(Y) \longrightarrow 0 \\ & & \downarrow \pi_X & & \downarrow [f \circ \pi_X, \alpha] & & \downarrow \pi_Y \\ 0 & \longrightarrow & X & \xrightarrow{f} & E & \xrightarrow{g} & Y \longrightarrow 0 \\ & & \downarrow j_X & & \downarrow \begin{bmatrix} \beta \\ j_Y \circ g \end{bmatrix} & & \downarrow j_Y \\ 0 & \longrightarrow & \mathfrak{I}(X) & \xrightarrow{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} & \mathfrak{I}(X) \oplus \mathfrak{I}(Y) & \xrightarrow{[0 \ 1]} & \mathfrak{I}(Y) \longrightarrow 0 \end{array}$$

Therefore

$$\begin{aligned} \bar{u}_E = \bar{u}_{E(\vec{x}_1)} &= \left[\begin{array}{c|c} \beta & \\ \hline j_Y \circ g & \end{array} \right] \circ \left[\begin{array}{c|c} f \circ \pi_X & \alpha \\ \hline \end{array} \right] = \left[\begin{array}{c|c} \beta \circ f \circ \pi_X & \beta \circ \alpha \\ \hline j_Y \circ g \circ f \circ \pi_X & j_Y \circ g \circ \alpha \end{array} \right] \\ &= \left[\begin{array}{c|c} j_X \circ \pi_X & \beta \circ \alpha \\ \hline 0 & j_Y \circ \pi_Y \end{array} \right] = \left[\begin{array}{c|c} \bar{u}_X(\vec{x}_1) & \beta \circ \alpha \\ \hline 0 & \bar{u}_Y(\vec{x}_1) \end{array} \right] = \left[\begin{array}{c|c} \bar{u}_X & \beta \circ \alpha \\ \hline 0 & \bar{u}_Y \end{array} \right]. \blacksquare \end{aligned}$$

From now on, we restrict to weighted projective lines \mathbb{X} of domestic type. For each of the weight types $(2, 2, n)$, $(2, 3, 3)$, $(2, 3, 4)$ and $(2, 3, 5)$ we determine in Propositions 3.6–3.9 the projective covers of indecomposable vector bundles, and thus by Theorem 3.3 also their injective hulls. The following propositions list from each Auslander–Reiten orbit of indecomposables of rank $r \geq 2$ a particular member, say E , and represent its projective cover $\mathfrak{P}(E) = \bigoplus_{i=1}^{2r} \mathcal{O}(\vec{y}_i)$ by the sequence $\vec{y}_1, \dots, \vec{y}_{2r}$ (including multiplicities). For the switch $\mathfrak{J}(E) = \mathfrak{P}(E)(\vec{x}_1)$ from projective covers to injective hulls we refer to the interpretation of the degree shift by \vec{x}_1 given in Remark 2.3.

Case $(2, 2, n)$. In this case the Auslander–Reiten quiver of $\text{vect}(\mathbb{X})$ has shape $\mathbb{Z}\widetilde{\mathbb{D}}_{n+2}$. Note that we need to distinguish the cases of n even (resp. n odd): For a given integral slope, there are exactly four (resp. two) line bundles if n is even (resp. if n is odd).



Here, the symbol \bullet (resp. \circ) marks the position of a vector bundle of rank 2 (resp. of a line bundle). Specifically, for key values of \vec{x} in \mathbb{L} , the position of the line bundle $\mathcal{O}(\vec{x})$ is indicated by the bracket symbol (\vec{x}) . Moreover, we have marked the position of $n - 1$ vector bundles E_0, \dots, E_{n-2} of rank two, one for each τ -orbit.


PROPOSITION 3.6. *We assume weight type $(2, 2, n)$ and refer to the notation in the above figure. Each indecomposable vector bundle that is not a*

line bundle has rank two. It lies in the τ -orbit of exactly one of the extension bundles E_0, \dots, E_{n-2} , where E_i is determined by the pair $(\mathcal{O}, i\vec{x}_3)$. Moreover,

$$\mathfrak{B}(E_i) = \mathcal{O}(\vec{\omega}) \oplus \mathcal{O}(i\vec{x}_3 - \vec{x}_1) \oplus \mathcal{O}(i\vec{x}_3 - \vec{x}_2) \oplus \mathcal{O}(-\vec{x}_3).$$

Proof. There are non-split exact sequences $0 \rightarrow \mathcal{O}(\vec{\omega}) \rightarrow E_i \rightarrow \mathcal{O}(i\vec{x}_3) \rightarrow 0$. Therefore E_i is isomorphic to the extension bundle $E\langle i\vec{x}_3 \rangle$, and the claim concerning projective covers follows from Theorem 3.2. ■

Case (2, 3, 3). In this case, we have three τ -orbits of line bundles, and the Auslander–Reiten quiver of $\text{vect}(\mathbb{X})$ has shape $\mathbb{Z}\tilde{\mathbb{E}}_6$:



For each τ -orbit of line bundles, we select one member, in bracket notation the line bundles \mathcal{O} , $\mathcal{O}(\vec{x}_2)$ and $\mathcal{O}(\vec{x}_3)$. Also, as indicated in the figure, for each of the four remaining τ -orbits, we select one member, resulting in three rank-two bundles E_2 , F_2 and G_2 and one bundle E_3 of rank three.

PROPOSITION 3.7. *We assume weight type (2, 3, 3) and refer to the notation in the above figure. Then each indecomposable bundle of rank at least two lies in the τ -orbit of exactly one of the vector bundles E_2 , F_2 , G_2 and E_3 , where the subscript indicates rank. The projective covers of these vector bundles are given in the table below:*

Vector bundle	Projective cover
E_2	$\vec{0}, -\vec{\omega} - \vec{x}_1, -\vec{\omega} - \vec{x}_2, -\vec{\omega} - \vec{x}_3$
F_2	$\vec{x}_2 - \vec{x}_3, -\vec{x}_2, \vec{x}_2 - \vec{x}_1, \vec{\omega}$
G_2	$2\vec{x}_2 - 2\vec{x}_3, -\vec{x}_3, \vec{\omega}, \vec{x}_3 - \vec{x}_1$
E_3	$\vec{\omega}, 2\vec{\omega}, \vec{x}_3 + 3\vec{\omega}, \vec{x}_3 + 4\vec{\omega}, \vec{x}_2 + 3\vec{\omega}, \vec{x}_2 + 4\vec{\omega}$

Proof. With the generator $u := \vec{x}_2 - \vec{x}_3$ of the torsion group $t\mathbb{L}$ of \mathbb{L} , the extension term H_i of the almost-split sequence $0 \rightarrow \mathcal{O}(i\vec{u}) \rightarrow H_i \rightarrow \mathcal{O}(i\vec{u} - \vec{\omega}) \rightarrow 0$ equals E_2 , F_2 or G_2 for $i = 0, 1$ or 2 , respectively. By means of Theorem 3.2, the claim on their projective covers follows. It thus remains

to determine the projective cover for E_3 . We note that each distinguished exact sequence with middle term of rank three necessarily splits. Hence for vector bundles of rank 3 it is not possible to use the horse-shoe argument from Lemma 3.4 in order to reduce the determination of projective covers to smaller rank. We thus need to give a direct argument:

Setting $F := E_2(\vec{\omega})$, we obtain short exact sequences

$$0 \rightarrow F \xrightarrow{i_{E_3}} E_3 \xrightarrow{\pi_{E_3}} \mathcal{O}(-\vec{\omega}) \rightarrow 0, \quad 0 \rightarrow \mathcal{O}(\vec{\omega}) \xrightarrow{i_F} F \xrightarrow{\pi_F} \mathcal{O} \rightarrow 0.$$

For $i = 1, 2, 3$ there are non-zero maps $x_i^{(-\vec{\omega})} : \mathcal{O}(-\vec{\omega} - \vec{x}_i) \rightarrow \mathcal{O}(-\vec{\omega})$, $x_i : \mathcal{O}(-\vec{x}_i) \rightarrow \mathcal{O}$. Because $\text{Ext}_{\mathbb{X}}^1(\mathcal{O}(-\vec{\omega} - \vec{x}_i), \mathcal{O}(\vec{\omega})) = 0 = \text{Ext}_{\mathbb{X}}^1(\mathcal{O}(-\vec{\omega} - \vec{x}_i), \mathcal{O})$ there are maps $y_i^{(-\vec{\omega})} : \mathcal{O}(-\vec{\omega} - \vec{x}_i) \rightarrow E_3$ such that $\pi_{E_3} \circ y_i^{(-\vec{\omega})} = x_i^{(-\vec{\omega})}$ for each $i = 1, 2, 3$. Since further $\text{Ext}_{\mathbb{X}}^1(\mathcal{O}(-\vec{x}_i), \mathcal{O}) = 0$, there are maps $y_i : \mathcal{O}(-\vec{x}_i) \rightarrow F$ such that $\pi_F \circ y_i = x_i$. We now show that each map $t : L \rightarrow E_3$, with L a line bundle, factors through $\pi = (i_{E_3} \circ i_F, (i_{E_3} \circ y_i), (y_i^{(-\vec{\omega})}))_{i=1,2,3}$.


We can assume that $\pi_{E_3} \circ t : L \rightarrow \mathcal{O}(-\vec{\omega})$ is not an isomorphism. Then $\pi_{E_3} \circ t = \sum_{i=1}^3 x_i^{(-\vec{\omega})} \circ t_i$, where $t_i : L \rightarrow \mathcal{O}(-\vec{x}_i \vec{\omega})$. Hence $\pi_{E_3} \circ t = \sum_{i=1}^3 x_i^{(-\vec{\omega})} \circ t_i = \sum_{i=1}^3 \pi_{E_3} \circ y_i^{(-\vec{\omega})} \circ t_i$, so $\pi_{E_3}(t - \sum_{i=1}^3 y_i^{(-\vec{\omega})} \circ t_i) = 0$. Therefore there is a map $g : L \rightarrow F$ such that $i_{E_3} \circ g = t - \sum_{i=1}^3 y_i^{(-\vec{\omega})} \circ t_i$. Again, $\pi_F \circ g$ is not an isomorphism, so $\pi_F \circ g = \sum_{i=1}^3 x_i \circ g_i$ for some $g_i : L \rightarrow \mathcal{O}(-\vec{x}_i)$. Then $\pi_F(g - \sum_{i=1}^3 y_i \circ g_i) = 0$. Hence there is a map $h : L \rightarrow \mathcal{O}(\vec{\omega})$ such that $i_F \circ h = g - \sum_{i=1}^3 y_i \circ g_i$. Applying i_{E_3} to this equality we obtain

$$t = i_{E_3} \circ i_F \circ h + \sum_{i=1}^3 [y_i^{(-\vec{\omega})} \circ t_i + (i_{E_3} \circ y_i) \circ g_i].$$

Thus t factors through $\pi = (i_{E_3} \circ i_F, (i_{E_3} \circ y_i), (y_i^{(-\vec{\omega})}))_{i=1,2,3}$.

Moreover, since $\text{Hom}_{\mathbb{X}}(\mathcal{O}(-\vec{x}_1), \mathcal{O}(-\vec{\omega} - \vec{x}_2))$ is non-zero and the spaces $\text{Hom}_{\mathbb{X}}(\mathcal{O}(-\vec{\omega} - \vec{x}_2), E_3)$ and $\text{Hom}_{\mathbb{X}}(\mathcal{O}(-\vec{x}_1), E_3)$ are one-dimensional, we can factor $y_1 : \mathcal{O}(-\vec{x}_1) \rightarrow E_3$ through $y_2^{(-\vec{\omega})} : \mathcal{O}(-\vec{\omega} - \vec{x}_2) \rightarrow E_3$. It is easy to see that the line bundles $\mathcal{O}(\vec{x}_3 + \vec{\omega})$, $\mathcal{O}(\vec{x}_2 + \vec{\omega})$, $\mathcal{O}(\vec{x}_1 + 2\vec{\omega})$, $\mathcal{O}(\vec{x}_3)$, $\mathcal{O}(\vec{x}_2)$, $\mathcal{O}(\vec{x}_1 + \vec{\omega})$ are Hom-orthogonal, which implies minimality of $\mathfrak{P}(E_3)$. Alternatively, minimality can be deduced from Theorem 3.3. ■

Case (2, 3, 4). In this case the Auslander–Reiten quiver of $\text{vect}(\mathbb{X})$ has shape $\mathbb{Z}\tilde{\mathbb{E}}_7$. It contains two τ -orbits of line bundles, three τ -orbits of rank-two bundles, two τ -orbits of rank-three bundles and one τ -orbit of rank-four bundles. We choose, in addition to the structure sheaf, one member of each τ -orbit, as indicated in the figure, and also mark the position of τG_2 :




PROPOSITION 3.8. *We assume weight type $(2, 3, 4)$ and refer to the notation in the above figure. Then each indecomposable vector bundle of rank at least two lies in the τ -orbit of exactly one of the vector bundles E_i for $i = 2, 3, 4$, $F_j = E_j(\vec{x}_1 - 2\vec{x}_3)$ for $j = 2, 3$, and $\tau_{\mathbb{X}}G_2$, where the subscript indicates rank. The projective covers of these vector bundles are given in the table below:*

Vector bundle	Projective cover
E_2	$\vec{0}, -\vec{\omega} - \vec{x}_1, -\vec{\omega} - \vec{x}_2, -\vec{\omega} - \vec{x}_3$
F_2	$\vec{\omega}, \vec{x}_2 - \vec{x}_1, -\vec{x}_2, \vec{x}_2 - \vec{x}_3$
$\tau_{\mathbb{X}}G_2$	$\vec{x}_2 - 2\vec{x}_3, -\vec{x}_2, \vec{x}_1 - 2\vec{x}_2, \vec{\omega} - \vec{x}_3$
E_3	$\vec{\omega}, \vec{x}_3 - \vec{x}_1, -\vec{x}_2, \vec{x}_2 - \vec{x}_1, -\vec{x}_3, \vec{x}_2 - \vec{x}_3$
F_3	$2\vec{x}_3 - \vec{x}_2, -\vec{x}_3, \vec{\omega} - \vec{x}_3, \vec{x}_2 - 2\vec{x}_3, \vec{x}_1 - 3\vec{x}_3, \vec{x}_1 - 2\vec{x}_2$
E_4	$\vec{x}_2 - 2\vec{x}_3, -\vec{x}_2, \vec{x}_2 - \vec{x}_1, \vec{\omega} - \vec{x}_3, \vec{\omega}, \vec{x}_3 - \vec{x}_1, \vec{x}_3 - \vec{x}_2, -\vec{x}_3$

Proof. Since $F_i = E_i(\vec{x}_1 - 2\vec{x}_3)$ for $i = 2, 3$ and $\mathfrak{P}(E(\vec{x})) = \mathfrak{P}(E)(\vec{x})$, it suffices to determine the projective covers of E_2, E_3, E_4 and of $\tau_{\mathbb{X}}G_2$. For the rank two bundles E_2 and $\tau_{\mathbb{X}}G_2$ this is an application of Theorem 3.2. For E_3 , the proof is similar to the proof of Proposition 3.7. It thus remains to determine the projective cover of E_4 by using Lemma 3.4. For this we consider the almost split sequence $0 \rightarrow \tau_{\mathbb{X}}G_2 \rightarrow E_4 \rightarrow G_2 \rightarrow 0$. If L is a line bundle summand of $\mathfrak{P}(G_2)$, then, by stability, $\mu L < \mu G_2$, and $\text{Ext}_{\mathbb{X}}^1(L, \tau_{\mathbb{X}}G_2) = D\text{Hom}_{\mathbb{X}}(G_2, L) = 0$. Similarly, if L' is a line bundle summand of $\mathfrak{P}(\tau_{\mathbb{X}}G_2)$, then $\text{Ext}_{\mathbb{X}}^1(G_2, L') = 0$. Therefore, the above sequence satisfies the assumptions of Lemma 3.4, hence $\mathfrak{P}(E_4) = \mathfrak{P}(\tau_{\mathbb{X}}G_2) \oplus \mathfrak{P}(G_2)$, as claimed. ■

Case $(2, 3, 5)$. In this case the Auslander–Reiten quiver of $\text{vect}(\mathbb{X})$ has the form $\mathbb{Z}\mathbb{E}_8$. We have just one τ -orbit of rank r for $r = 1, 5, 6$, and two

τ -orbits of rank r for each $r = 2, 3, 4$.



The marked region is a fundamental domain with respect to the Auslander–Reiten translation.

PROPOSITION 3.9. *We assume weight type $(2, 3, 5)$ and refer to the notation in the above figure. Then each indecomposable bundle of rank ≥ 2 lies in the Auslander–Reiten orbit of exactly one of the vector bundles E_i , F_j and G_l , having rank $i = 2, 3, 4, 5, 6$ (resp. $j = 2, 4$, $l = 3$). Moreover, the projective covers are given in the table below:*

Vector bundle	Projective cover
E_2	$\vec{0}, \vec{x}_3 - 2\vec{x}_2, \vec{x}_3 - \vec{x}_1, \vec{x}_2 - \vec{x}_1$
F_2	$\vec{x}_1 - 3\vec{x}_3, \vec{\omega} - \vec{x}_2, -\vec{x}_3, \vec{x}_2 - 3\vec{x}_3$
E_3	$\vec{\omega}, \vec{x}_2 - \vec{x}_1, -\vec{x}_3, \vec{x}_3 - \vec{x}_1, -\vec{x}_2, \vec{x}_3 - 2\vec{x}_2$
G_3	$\vec{x}_1 - 3\vec{x}_3, \vec{x}_2 - \vec{x}_1, \vec{\omega} - \vec{x}_3, \vec{x}_2 - 3\vec{x}_3, -\vec{x}_2, -2\vec{x}_3$
E_4	$\vec{x}_2 - \vec{x}_1, \vec{\omega} - 2\vec{x}_3, \vec{x}_2 - 3\vec{x}_3, -\vec{x}_2, \vec{\omega}, \vec{x}_3 - \vec{x}_1, \vec{x}_3 - \vec{x}_2, -\vec{x}_3$
F_4	$\vec{x}_3 - \vec{x}_2, -2\vec{x}_3, -\vec{x}_3, \vec{x}_3 - \vec{x}_1, \vec{x}_1 - 3\vec{x}_3, \vec{\omega} - \vec{x}_2, -\vec{x}_3, \vec{x}_2 - 3\vec{x}_3$
E_5	$\vec{x}_3 - \vec{x}_2, -\vec{x}_3, \vec{x}_2 - 3\vec{x}_3, \vec{x}_3 - \vec{x}_1, \vec{\omega} - \vec{x}_2, \vec{\omega} - 2\vec{x}_3, \vec{x}_2 - 2\vec{x}_3, -\vec{x}_2, \vec{x}_2 - \vec{x}_1, \vec{\omega} - \vec{x}_3$
E_6	$\vec{x}_1 - 3\vec{x}_3, \vec{x}_2 - \vec{x}_1, \vec{\omega} - \vec{x}_3, \vec{x}_2 - 3\vec{x}_3, -\vec{x}_2, -2\vec{x}_3, \vec{x}_2 - 2\vec{x}_3, \vec{x}_3 - \vec{x}_2, -\vec{x}_3, \vec{\omega} - \vec{x}_3, \vec{x}_3 - \vec{x}_1, \vec{\omega} - \vec{x}_2$

Proof. In the case of vector bundles of rank 2 and 3 the proof is similar to that of Proposition 3.7. For the remaining cases we will use Lemma 3.4.

In the case of E_4 we consider the exact sequence $0 \rightarrow \tau_{\mathbb{X}}^2 F_2 \rightarrow E_4 \rightarrow \tau_{\mathbb{X}}^{-2} F_2 \rightarrow 0$. It is easy to see that it satisfies the assumptions of Lemma 3.4, i.e.

$$\begin{aligned}\mathrm{Ext}_{\mathbb{X}}^1(\mathfrak{P}(\tau_{\mathbb{X}}^{-2}F_2), \tau_{\mathbb{X}}^2F_2) &= D\mathrm{Hom}_{\mathbb{X}}(\tau_{\mathbb{X}}F_2, \mathfrak{P}(F_2)) = 0, \\ \mathrm{Ext}_{\mathbb{X}}^1(\tau_{\mathbb{X}}^{-2}F_2, \mathfrak{J}(\tau_{\mathbb{X}}^2F_2)) &= D\mathrm{Hom}_{\mathbb{X}}(\tau_{\mathbb{X}}\mathfrak{J}(F_2), F_2) = 0.\end{aligned}$$

Hence $\mathfrak{P}(E_4) = \mathfrak{P}(\tau_{\mathbb{X}}^2F_2) \oplus \mathfrak{P}(\tau_{\mathbb{X}}^{-2}F_2)$, so the results follow from Theorem 3.2 applied to the extension bundle $\tau_{\mathbb{X}}^2F_2$ (resp. $\tau_{\mathbb{X}}^{-2}F_2$), which is determined by the pair $(\mathcal{O}(\vec{x}_3 - \vec{x}_2), \vec{x}_3)$ (resp. (\mathcal{O}, \vec{x}_3)).

For F_4 , E_5 and E_6 , we use the exact sequences $0 \rightarrow \tau_{\mathbb{X}}F_2 \rightarrow F_4 \rightarrow F_2 \rightarrow 0$, $0 \rightarrow \tau_{\mathbb{X}}G_3 \rightarrow E_5 \rightarrow \tau_{\mathbb{X}}^{-1}F_2 \rightarrow 0$ and $0 \rightarrow G_3 \rightarrow E_6 \rightarrow \tau_{\mathbb{X}}^{-1}G_3 \rightarrow 0$, respectively. It is straightforward to check that these satisfy the conditions of Lemma 3.4. ■

REMARK 3.10. Later, when calculating matrix factorizations for E_6 , we will use two different exact sequences, namely $0 \rightarrow G_3 \rightarrow E_6 \rightarrow \tau_{\mathbb{X}}^{-1}G_3 \rightarrow 0$ and $0 \rightarrow \tau_{\mathbb{X}}F_4 \rightarrow E_6 \rightarrow \tau_{\mathbb{X}}^{-1}F_2 \rightarrow 0$. While both yield the same projective cover, we will obtain different matrix factorizations, because the two procedures yield matrix factorizations with a different number of zero entries.

We conclude this section with the observation that indecomposable vector bundles are uniquely determined by their projective covers, provided \mathbb{X} is domestic. This result turns out to be central to determining matrix factorizations for indecomposable vector bundles.

PROPOSITION 3.11. *We assume a domestic weight triple $(2, a, b)$. Let E and F be indecomposable vector bundles. Then E and F are isomorphic if and only if their projective covers $\mathfrak{P}(E)$ and $\mathfrak{P}(F)$ are isomorphic.*

Proof. We may assume that $\mathfrak{P}(E) = \mathfrak{P}(F)$ where E and F have rank at least two.

The first part of the proof holds for arbitrary weight triples $(2, a, b)$: From the (distinguished) exact sequence $0 \rightarrow E(-\vec{x}_1) \rightarrow \mathfrak{P}(E) \rightarrow E \rightarrow 0$, using (2.4) we obtain $\det(\mathfrak{P}(E)) = 2 \det(E) + \vec{c}$, and moreover $\mathrm{rk}(\mathfrak{P}(E)) = 2 \mathrm{rk}(E)$. Therefore $\mathfrak{P}(E)$ determines the determinant, degree, rank and slope of E . In particular, $\mathfrak{P}(E) = \mathfrak{P}(F)$ implies that E and F have the same rank and the same slope.

Next, we establish the claim separately for the domestic weight triples $(2, 2, n)$, $(2, 3, 3)$, $(2, 3, 4)$ and $(2, 3, 5)$.

CASE $(2, 2, n)$: We refer to the notation of Proposition 3.6. By an appropriate τ -shift, we may assume that the bundles E_0, E_2, E_4, \dots (resp. the bundles E_1, E_3, E_5, \dots) have the same slope. One checks that the bundles in each of the two families have distinct systems of line bundle summands in the projective cover having maximal slope.

CASE $(2, 3, 3)$: With the notation of Proposition 3.7, the rank-two bundles E_2 , F_2 and G_2 have the same slope, but different line bundle summands of their projective cover with maximal slope, namely \mathcal{O} , $\mathcal{O}(\vec{x}_3 - \vec{x}_2)$

and $\mathcal{O}(2(\vec{x}_3 - \vec{x}_2))$, respectively. Since there is a unique τ -orbit of indecomposable rank-three bundles, each of these bundles is determined by its slope.

CASE (2, 3, 4): We refer to the notation of Proposition 3.8. The rank-two bundles E_2 and $E_2(\vec{x}_3 - 2\vec{x}_1)$ have the same half-integral slope and belong to different τ -orbits. They have different line bundle summands of maximal slope in their respective projective covers, namely \mathcal{O} and $\mathcal{O}(\vec{x}_3 - 2\vec{x}_1)$. The members from the third τ -orbit of rank-two bundles, in particular T , are distinguished from members from the other two τ -orbits by their slope which is integral. In a similar way, the bundles E_3 and $E_3(\vec{x}_3 - 2\vec{x}_1)$ have the same slope, they represent the τ -orbits of rank-three bundles, and have different line bundle summands of maximal slope in their projective covers, namely $\mathcal{O}(\vec{\omega})$ and $\mathcal{O}(\vec{\omega} + \vec{x}_3 - 2\vec{x}_1)$. Finally, there is just one τ -orbit of indecomposable rank-four bundles.

CASE (2, 3, 5): We refer to the notation of Proposition 3.9. Here, the claim reduces to showing that E_2, F_2 (analogously E_3, F_3 and E_4, F_4) can be distinguished in terms of their projective covers. To distinguish E_2 and F_2 we observe that \mathcal{O} and \mathcal{O} , respectively $\mathcal{O}(3\vec{\omega})$, are line bundle summands of their projective covers. Concerning E_3 and F_3 , the line bundle summands \mathcal{O} and $\mathcal{O}(4\vec{\omega})$ have maximal slopes in their respective projective covers. Finally, the Auslander–Reiten orbits of E_4 and F_4 are distinguished by their integral (resp. half-integral) slopes. ■

Assuming an arbitrary weight triple, a corresponding result holds true for indecomposable bundles of rank two. For the proof we refer to [LR].

PROPOSITION 3.12. *We assume that \mathbb{X} has triple weight type (a, b, c) . Then each indecomposable vector bundle E of rank two is uniquely determined by its projective cover $\mathfrak{P}(E)$. ■*

REMARK 3.13. Assuming a weighted projective line $\mathbb{X} = \mathbb{X}(a, b, c)$ of Euler characteristic $\chi_{\mathbb{X}} \leq 0$, that is, assuming \mathbb{X} of tubular or wild type, it is no longer true that each indecomposable vector bundle E is uniquely determined by $\mathfrak{P}(E)$. Suppose that the base field k is uncountable and, for simplicity, that \mathbb{X} has weight type $(2, a, b)$. By perpendicular calculus (see [GL91]) there exists a weighted projective line \mathbb{Y} of tubular type and a full embedding $\text{coh}(\mathbb{Y}) \hookrightarrow \text{coh}(\mathbb{X})$ that preserves the rank. From the tubular families in $\text{coh}(\mathbb{Y})$ we then deduce the existence of a one-parameter family (E_α) of indecomposable vector bundles over \mathbb{X} , all having the same positive rank r . This in turn implies that each projective hull $\mathfrak{P}(E_\alpha)$ has fixed rank $2r$. Since the grading group \mathbb{L} is countable, this leaves only countably many possibilities for the isomorphism classes of $\mathfrak{P}(E_\alpha)$, forcing many non-isomorphic E_α 's to have the same projective cover.

This leads to a modified question, to which the authors do not know the answer: *Assume that E and F are exceptional vector bundles with isomorphic projective covers $\mathfrak{P}(E)$ and $\mathfrak{P}(F)$. Does this imply that E and F are isomorphic?* In support for a positive answer, we mention that, for \mathbb{X} domestic, each indecomposable vector bundle is exceptional. Also, because of triple weight type, all extension bundles are exceptional. Moreover, since by a result of Hübner [Hüb96] (see [Mel04] for a proof), exceptional vector bundles are determined by their classes in the Grothendieck group $K_0\mathbb{X}$, there exist only countably many isoclasses of exceptional vector bundles, thus preventing the contradiction of the above argument.

4. Factorization frame attached to a vector bundle. We assume a domestic weight triple $(2, a, b) = (p_1, p_2, p_3)$. Then a matrix factorization for an indecomposable vector bundle E of rank r can be obtained from its minimal projective resolution by first determining its *factorization frame* consisting of a pair of $2r \times 2r$ -matrices (see Definition 4.4), and then adjusting the entries of the factorization frame by suitable scalars.

We now establish the key result of this section. For this, it is convenient to identify the Frobenius categories $\mathrm{CM}^{\mathbb{L}}(S)$ and $\mathrm{vect}(\mathbb{X})$ by means of sheafification $\mathrm{CM}^{\mathbb{L}}(S) \xrightarrow{\sim} \mathrm{vect}(\mathbb{X})$, $M \mapsto \tilde{M}$. In the same context, by a matrix factorization (u, v) attached to a vector bundle E without line bundle summands we mean a matrix factorization of $f = x_1^{p_1} + x_2^{p_2} + x_3^{p_3}$ over $T = k[x_1, x_2, x_3]$ attached to the Cohen–Macaulay module M corresponding to E . We are going to compare the rows of the commutative diagram

$$(4.1) \quad \begin{array}{ccccccccc} \cdots & \longrightarrow & P_0(-\vec{c}) & \xrightarrow{v} & P_1 & \xrightarrow{u} & P_0 & \xrightarrow{\pi} & M & \longrightarrow & 0 \\ & & \nu \downarrow & & \nu \downarrow & & \nu \downarrow & & \parallel & & \\ \cdots & \longrightarrow & \bar{P}_0(-\vec{c}) & \xrightarrow{\bar{v}} & \bar{P}_1 & \xrightarrow{\bar{u}} & \bar{P}_0 & \xrightarrow{\bar{\pi}} & M & \longrightarrow & 0 \end{array}$$

where the upper row is a T -matrix factorization of f for M , π is a T -projective cover of M , and the lower row is a minimal S -projective resolution of M . The vertical maps and the bar notation stand for reduction modulo (f) .

In the above setting, we fix decompositions of P_0 and P_1 into indecomposable T -projectives, and consider the decompositions for the T -projectives $P_0(-n\vec{c})$, induced by degree shift, and the corresponding decompositions of the S -projectives $\bar{P}_0(-n\vec{c})$ and $\bar{P}_1(-n\vec{c})$, induced by reduction modulo (f) . We then say that we have chosen *compatible decompositions* for (4.1). To achieve such decompositions, we may alternatively start with decompositions of \bar{P}_0 and \bar{P}_1 , and then extend them to the remaining members of (4.1) by degree shift and by taking T -projective covers.

We say that an element $x = x_1^{l_1} x_2^{l_2} x_3^{l_3}$, viewed as a member of T or S , is a *monomial with small exponents* if $0 \leq l_i \leq p_i - 1$ for $i = 1, 2, 3$ and $\sum_{i=1}^3 l_i > 0$. In particular, x belongs to the graded maximal ideal of T (respectively S).

THEOREM 4.1. *We assume the above setting (4.1) for a weighted projective line \mathbb{X} of domestic type, where M is an indecomposable \mathbb{L} -graded Cohen–Macaulay module of rank at least two, attached to the (indecomposable) vector bundle E on \mathbb{X} .*

Let \vec{y}_0, \vec{y}_1 be members of \mathbb{L} such that $T(\vec{y}_i)$ is a T -direct summand of P_i ($i = 0, 1$) and, accordingly, $S(\vec{y}_i)$ is an S -direct summand of \bar{P}_i . Then reduction modulo (f) induces an isomorphism

$$(4.2) \quad T_{\vec{y}_0 - \vec{y}_1} = \mathrm{Hom}_T(T(\vec{y}_1), T(\vec{y}_0)) \xrightarrow{\cong} \mathrm{Hom}_S(S(\vec{y}_1), S(\vec{y}_0)) = S_{\vec{y}_0 - \vec{y}_1}.$$

Moreover, the Hom-spaces from (4.2) are either both zero, and then $\vec{y}_1 \not\leq \vec{y}_0$, or else $0 < \delta(\vec{y}_0 - \vec{y}_1) < \delta(\vec{c})$, and then $T_{\vec{y}_0 - \vec{y}_1} = kx$ (and also $S_{\vec{y}_0 - \vec{y}_1} = kx$) for a monomial x with small exponents.

Proof. Since f belongs to the graded maximal ideal of S , a finitely generated graded T -module M is zero if and only if M/fM is zero. In particular, $\mathrm{Hom}_T(T(\vec{y}_1), T(\vec{y}_0)) = 0$ if and only if $\mathrm{Hom}_S(S(\vec{y}_1), S(\vec{y}_0)) = 0$.

Next, we switch to the context of vector bundles, and use the existence of (distinguished) exact sequences $0 \rightarrow E(-\vec{x}_1) \rightarrow \bar{P}_0 \rightarrow E \rightarrow 0$ and $0 \rightarrow E(-\vec{c}) \rightarrow \bar{P}_1 \rightarrow E(-\vec{x}_1) \rightarrow 0$, where the \bar{P}_0 and \bar{P}_1 are projective in the Frobenius category $\mathrm{vect}(\mathbb{X})$. Assume that $\mathrm{Hom}_{\mathbb{X}}(\mathcal{O}(\vec{y}_1), \mathcal{O}(\vec{y}_0)) \neq 0$. We are going to show that $\mathrm{Hom}_{\mathbb{X}}(\mathcal{O}(\vec{y}_1), \mathcal{O}(\vec{y}_2)) = kx$, where x is a small monomial: If E has slope $\mu E = q$, then $\mu E(-\vec{x}_1) = q - \delta(\vec{x}_1)$ and $\mu E(-\vec{c}) = q - \delta(\vec{c})$. Because \mathbb{X} is domestic, all indecomposable vector bundles are stable by [GL87, Proposition 5.5]. Since \bar{P}_1 is the injective hull of $E(-\vec{c})$, we have $\mathrm{Hom}_{\mathbb{X}}(E(-\vec{c}), \mathcal{O}(\vec{y}_1)) \neq 0$, hence $q - \delta(\vec{c}) < \delta(\vec{y}_1)$. Since \bar{P}_1 is the projective cover of $E(-\vec{x}_1)$, we have $\mathrm{Hom}_{\mathbb{X}}(\mathcal{O}(\vec{y}_1), E(-\vec{x}_1)) \neq 0$, hence $\delta(\vec{y}_1) < q - \delta(\vec{x}_1)$. Similarly, because \bar{P}_0 is the injective hull of $E(-\vec{x}_1)$ and also the projective cover of E , we obtain $q - \delta(\vec{x}_1) < \delta(\vec{y}_0) < q$. Putting things together, we finally get

$$q - \delta(\vec{c}) < \delta(\vec{y}_1) < q - \delta(\vec{x}_1) < \delta(\vec{y}_0) < q,$$

and in particular $\delta(\vec{y}_0 - \vec{y}_1) < \delta(\vec{c})$. Since $\mathrm{Hom}_{\mathbb{X}}(\mathcal{O}(\vec{y}_1), \mathcal{O}(\vec{y}_0)) = S_{\vec{y}_1 - \vec{y}_0}$ we get $0 < \delta(\vec{y}_1 - \vec{y}_0) < \delta(\vec{c})$. We set $\vec{u} = \vec{y}_0 - \vec{y}_1$. Since $S_{\vec{y}_0 - \vec{y}_1} \neq 0$ by assumption, we get $0 < \vec{u}$ and $0 < \delta(\vec{u}) < \delta(\vec{c})$. Note that $\vec{u} = 0$ is not possible, because $\delta(\vec{u}) > 0$. Writing $\vec{y}_0 - \vec{y}_1$ in normal form $\vec{y}_0 - \vec{y}_1 = l_1 \vec{x}_1 + l_2 \vec{x}_2 + l_3 \vec{x}_3 + l \vec{c}$ with $0 \leq l_i \leq p_i - 1$, we obtain $l \geq 0$ from $\vec{y}_0 - \vec{y}_1 > 0$. Assuming that $l \geq 1$ then yields $\delta(\vec{y}_0 - \vec{y}_1) \geq \delta(\vec{c})$, which is impossible. Thus $l = 0$ and $S_{\vec{y}_0 - \vec{y}_1} = k\vec{x}_1^{l_1} \vec{x}_2^{l_2} \vec{x}_3^{l_3}$, establishing the last

assertion. From this it finally follows that the map from (4.2) is an isomorphism. ■

For the corollary below, we keep the notation and assumptions of Theorem 4.1.

COROLLARY 4.2. *We assume compatible decompositions for the members of (4.1). Then the T -matrix factorization (u, v) of f , associated to E , and the S -minimal projective resolution (\bar{u}, \bar{v}) are represented by the ‘same’ matrix pair (U, V) , whose entries are scalar multiples of monomials with small exponents, interpreted as elements of T (respectively of S). ■*

The scalar factors from Corollary 4.2, and hence the matrices (U, V) , are usually difficult to determine. We thus introduce an intermediate concept, called a *factorization frame* for E . By Theorem 4.1 or Corollary 4.2, factorization frames always exist for indecomposable bundles, provided we deal with domestic weight triples. But factorization frames may also exist in other situations. In particular, extension bundles admit factorization frames, without any restriction on the weight triple (p_1, p_2, p_3) .

Assuming an arbitrary weight triple (a, b, c) , Theorem 3.2 provides an explicit projective cover for extension bundles, and thus for indecomposable vector bundles of rank two. Hence a result very close to Theorem 4.1 can be shown for indecomposable Cohen–Macaulay modules of rank two for arbitrary weight triples (a, b, c) , by just following the lines of the proof for Theorem 4.1:

THEOREM 4.3. *We assume the above setting (4.1) for a weighted projective line \mathbb{X} of type (a, b, c) , where M is an indecomposable \mathbb{L} -graded Cohen–Macaulay module of rank two.*

Let \vec{y}_0, \vec{y}_1 be members of \mathbb{L} such that $T(\vec{y}_i)$ is a T -direct summand of P_i ($i = 0, 1$) and, accordingly, $S(\vec{y}_i)$ is an S -direct summand of \bar{P}_i . Then reduction modulo (f) induces an isomorphism

$$(4.3) \quad T_{\vec{y}_0-\vec{y}_1} = \mathrm{Hom}_T(T(\vec{y}_1), T(\vec{y}_0)) \xrightarrow{\cong} \mathrm{Hom}_S(S(\vec{y}_1), S(\vec{y}_0)) = S_{\vec{y}_0-\vec{y}_1}.$$

Moreover, the Hom-spaces from (4.3) are either both zero, in which case $\vec{y}_1 \not\leq \vec{y}_0$, or else $0 < \delta(\vec{y}_0 - \vec{y}_1) < \delta(\vec{c})$, in which case $T_{\vec{y}_0-\vec{y}_1} = kx$ (and also $S_{\vec{y}_0-\vec{y}_1} = kx$) for a monomial x with small exponents.

DEFINITION 4.4. We assume a domestic weight triple. A *factorization frame* (U, V) for an indecomposable vector bundle E of rank $r \geq 2$ is a pair of $2r \times 2r$ -matrices over T , obtained by Theorem 4.3 from line bundle decompositions

$$(4.4) \quad \bar{P}_0 = \bigoplus_{j=1}^{2r} \mathcal{O}(\vec{z}_j), \quad \bar{P}_1 = \bigoplus_{i=1}^{2r} \mathcal{O}(\vec{y}_i), \quad \bar{P}_0(-\vec{c}) = \bigoplus_{j=1}^{2r} \mathcal{O}(\vec{z}_j - \vec{c})$$

of the projectives from a minimal projective resolution of E , as follows: The (i, j) -entry of U is defined by:

- (a) If $\mathrm{Hom}_{\mathbb{X}}(\mathcal{O}(\vec{y}_i), \mathcal{O}(\vec{z}_j)) = 0$, then the entry is zero.
- (b) Otherwise, $\mathrm{Hom}_{\mathbb{X}}(\mathcal{O}(\vec{y}_i), \mathcal{O}(\vec{z}_j)) = S_{\vec{z}_j - \vec{y}_i} = kx_1^{l_1}x_2^{l_2}x_3^{l_3}$ with $0 \leq l_i \leq (p_i - 1)$. Then the (i, j) -entry is given by the monomial $x_1^{l_1}x_2^{l_2}x_3^{l_3}$.

The matrix V is defined in a similar fashion from the decompositions of $P_0(-\vec{c})$ and P_1 .

By Definition 4.4, a factorization frame for E depends on the line bundle decompositions for the terms of a minimal projective resolution of E . However, assuming domestic type, for $\mathrm{rk}(E) \leq 5$ the factorization frame attached to E is unique by Proposition 4.6. We note also that usually a factorization frame (U, V) for E will not satisfy the matrix factorization property $UV = f\mathbf{1} = VU$. However, by Corollary 4.2, each factorization frame for E can be specialized to a matrix factorization for E :

LEMMA 4.5. *We assume domestic type and that E is indecomposable of rank at least two. Then each factorization frame (U, V) for E can be specialized to a T -matrix factorization for f , representing E , by modifying the entries of the factorization frame by (possibly zero) scalars in such a way that the resulting matrices u, v satisfy $uv = f\mathbf{1} = vu$. Conversely, each matrix factorization (u, v) for f , representing E , arises this way. ■*

Let E be an indecomposable vector bundle of rank ≥ 2 for domestic weight type. Our next result implies that, with the single exception of the members E from the single τ -orbit of indecomposable rank-six bundles for weight type $(2, 3, 5)$, the decomposition of the projective cover $\mathfrak{P}(E)$ into line bundles is multiplicity-free.

PROPOSITION 4.6. *Let \mathbb{X} be of domestic type and let E be an indecomposable vector bundle of rank $r \geq 2$, with projective cover $\mathfrak{P}(E) = \bigoplus_{i=1}^{2r} L_i$. If $\mathrm{rk}(E) \leq 5$, then L_1, \dots, L_{2r} are pairwise non-isomorphic line bundles. If $\mathrm{rk}(E) = 6$, and then \mathbb{X} of weight type $(2, 3, 5)$, there are exactly 11 non-isomorphic line bundles among L_1, \dots, L_{12} .*

Proof. Case-by-case analysis, based on Propositions 3.6–3.9. ■

Of course, not every matrix factorization (u, v) , obtained from a factorization frame (U, V) , attached to an indecomposable vector bundle E by specialization, will satisfy $\mathrm{cok}(u, v) = E$: see Example 4.12, where the matrices u and v contain too many zero entries. We thus investigate situations when the modified entries of a factorization frame are not allowed to be zero. For this the following general result from [KLM13, Prop. 3.8] will be useful. Here, we say that a map $h : E \rightarrow L'$ is a *component map* of the injective hull $j_E : E \rightarrow \mathfrak{J}(E)$ if L' is a line bundle and $E = L' \oplus E'$ where h is the

restriction of j_E to L' . Component maps of projective covers are defined in a similar way.

PROPOSITION 4.7. *Let \mathbb{X} be a weighted projective line of triple weight type. Let E be a vector bundle and L, L' be line bundles.*

- (i) *If $h : E \rightarrow L'$ is a component map of the injective hull $j_E : E \rightarrow \mathfrak{J}(E)$, then h is an epimorphism in $\text{coh}(\mathbb{X})$.*
- (ii) *If $l : L \rightarrow E$ is a component map of the projective cover $\pi_E : \mathfrak{P}(E) \rightarrow E$, then l is a monomorphism in $\text{coh}(\mathbb{X})$, and moreover the cokernel of l , formed in $\text{coh}(\mathbb{X})$, is a vector bundle.*

Proof. Property (i) immediately follows from the explicit description of the injective hull in Theorem 3.2, while the proof of (ii) uses the case-by-case description of projective covers from Section 3. ■

As an immediate consequence we get

COROLLARY 4.8. *Let \mathbb{X} be a weighted projective line of triple weight type. For an exceptional vector bundle E we have*

- (i) $\text{Ext}_{\mathbb{X}}^1(E, \mathfrak{J}(E)) = 0$,
- (ii) $\text{Ext}_{\mathbb{X}}^1(\mathfrak{P}(E), E) = 0$,
- (iii) $\text{Ext}_{\mathbb{X}}^1(\mathfrak{P}(E), \mathfrak{J}(E)) = 0$.

Proof. For (i), let L' be a line bundle summand of $\mathfrak{J}(E)$ and $v : E \rightarrow L'$ be a corresponding component map of the injective hull $j_E : E \rightarrow \mathfrak{J}(E)$. By Proposition 4.7, $v : E \rightarrow L'$ is an epimorphism. Thus, by heredity of $\text{coh}(\mathbb{X})$, the condition $\text{Ext}_{\mathbb{X}}^1(E, E) = 0$ implies that $\text{Ext}_{\mathbb{X}}^1(E, L') = 0$. This happens for each line bundle summand L' of $\mathfrak{J}(E)$, therefore $\text{Ext}_{\mathbb{X}}^1(E, \mathfrak{J}(E)) = 0$. The proof of (ii) is dual. Concerning (iii), we argue as before: Since v is an epimorphism, condition (ii) implies $\text{Ext}_{\mathbb{X}}^1(\mathfrak{P}(E), L') = 0$ for each line bundle summand L' of $\mathfrak{J}(E)$, and (iii) follows. ■

LEMMA 4.9. *Let \mathbb{X} be a weighted projective line of triple weight type and let E be an exceptional vector bundle. Let $\pi_E : \mathfrak{P}(E) \rightarrow E$ (respectively $j_E : E \rightarrow \mathfrak{J}(E)$) be the projective cover (respectively the injective hull) of E . Let L (respectively L') be a line bundle summand of the projective cover $\mathfrak{P}(E)$ (respectively the injective hull $\mathfrak{J}(E)$), and let $u : L \rightarrow E$ (respectively $v : E \rightarrow L'$) be a corresponding component map of $\pi_E : \mathfrak{P}(E) \rightarrow E$ (respectively $j_E : E \rightarrow \mathfrak{J}(E)$). Assume that $\text{Hom}_{\mathbb{X}}(L, E) = k$ and $\text{Hom}_{\mathbb{X}}(E, L') = k$. Then vu is the zero map if and only if $\text{Hom}_{\mathbb{X}}(L, L') = 0$.*

Proof. Assume, for contradiction, that $\text{Hom}_{\mathbb{X}}(L, L') \neq 0$ but $vu = 0$. By Proposition 4.7 the map $v : E \rightarrow L'$ is an epimorphism. By heredity of $\text{coh}(\mathbb{X})$, the condition $\text{Ext}_{\mathbb{X}}^1(E, E) = 0$ then implies that $\text{Ext}_{\mathbb{X}}^1(E, L') = 0$.

Next, we apply the functor $\mathrm{Hom}_{\mathbb{X}}(-, L')$ to the non-split exact sequence $(\star) 0 \rightarrow L \xrightarrow{u} E \xrightarrow{p} F \rightarrow 0$, yielding exactness of $\mathrm{Hom}_{\mathbb{X}}(E, L') \xrightarrow{-\circ u} \mathrm{Hom}_{\mathbb{X}}(L, L') \rightarrow \mathrm{Ext}_{\mathbb{X}}^1(F, L') \rightarrow \mathrm{Ext}_{\mathbb{X}}^1(E, L') = 0$. Because $\mathrm{Hom}_{\mathbb{X}}(E, L') = k$ and $v \circ u = 0$, the map $-\circ u : \mathrm{Hom}_{\mathbb{X}}(E, L') \rightarrow \mathrm{Hom}_{\mathbb{X}}(L, L')$ is zero. Therefore $\mathrm{Ext}_{\mathbb{X}}^1(F, L') \cong \mathrm{Hom}_{\mathbb{X}}(L, L') \neq 0$ by assumption. On the other hand, applying $\mathrm{Hom}_{\mathbb{X}}(-, E)$ to (\star) , we get exactness of $0 \rightarrow \mathrm{Hom}_{\mathbb{X}}(F, E) \rightarrow \mathrm{Hom}_{\mathbb{X}}(E, E) \rightarrow \mathrm{Hom}_{\mathbb{X}}(L, E) \rightarrow \mathrm{Ext}_{\mathbb{X}}^1(F, E) \rightarrow \mathrm{Ext}_{\mathbb{X}}^1(E, E) = 0$. Since, by assumption, $\mathrm{Hom}_{\mathbb{X}}(E, E) = k = \mathrm{Hom}_{\mathbb{X}}(L, E)$, we deduce that $\dim \mathrm{Hom}_{\mathbb{X}}(F, E) = \dim \mathrm{Ext}_{\mathbb{X}}^1(F, E)$, and we are going to show that both terms vanish.

Assuming $\mathrm{Hom}_{\mathbb{X}}(F, E) \neq 0$, we compose the epimorphism p with a non-zero map from F to E to obtain an endomorphism of E that is neither zero nor an isomorphism, in obvious contradiction to $\mathrm{Hom}(E, E) = k$. Hence $\mathrm{Hom}_{\mathbb{X}}(F, E) = 0$ and then $\mathrm{Ext}_{\mathbb{X}}^1(F, E) = 0$. Because $v : E \rightarrow L'$ is an epimorphism, the condition $\mathrm{Ext}_{\mathbb{X}}^1(F, E) = 0$ finally implies that $\mathrm{Ext}_{\mathbb{X}}^1(F, L') = 0$, contrary to what was established before. ■

The following consequence yields certain limitations for specializing factorization frames to matrix factorizations. We adhere to the notation of Definition 4.4 and further refer to Lemma 4.5.

COROLLARY 4.10. *Assume a matrix factorization (u, v) of f over T that is attached to E , is obtained from a factorization frame (U, V) for E by specialization. Assume, in particular, that $u_{ij} = \lambda_{ij}U_{ij}$. If $\mathrm{Hom}(\mathcal{O}(\vec{y}_i), E) = k = \mathrm{Hom}(E, \mathcal{O}(\vec{z}_j))$, then the scalar λ_{ij} must be non-zero. A similar result holds for the specialization of V to v .*

In the general situation the dimension of the homomorphism space between E and a line bundle summand of $\mathfrak{P}(E)$ or $\mathfrak{J}(E)$ can be greater than one. We have more precise information for indecomposable vector bundles of rank 2 and 3:

LEMMA 4.11.

- (a) *Assuming \mathbb{X} of arbitrary weight triple, let L (respectively L') be a direct summand of the projective cover (respectively the injective hull) of an extension bundle E . Then $\mathrm{Hom}_{\mathbb{X}}(L, E) = k$ (respectively $\mathrm{Hom}_{\mathbb{X}}(E, L') = k$).*
- (b) *Assuming \mathbb{X} of domestic weight type, let L (respectively L') be a direct summand of the projective cover (respectively the injective hull) of an indecomposable rank 3 bundle E . Then $\mathrm{Hom}_{\mathbb{X}}(L, E) = k$ (respectively $\mathrm{Hom}_{\mathbb{X}}(E, L') = k$).*

Proof. We prove (a); the proof of (b) is similar. Let E be an extension bundle on \mathbb{X} , thus E is the middle term of an exact sequence $0 \rightarrow \bar{L}(\vec{\omega}) \rightarrow E \rightarrow \bar{L}(\vec{x}) \rightarrow 0$ for some line bundle \bar{L} and an element $\vec{x} = l_1\vec{x}_1 + l_2\vec{x}_2 + l_3\vec{x}_3$

such that $0 \leq l_i \leq p_i - 2$ for $i = 1, 2, 3$. It follows from Theorem 3.2 that each direct summand L of $\mathfrak{P}(E)$ is $\bar{L}(\vec{\omega})$ or $\bar{L}(\vec{x} - (1 + l_i \vec{x}_i))$ for $i = 1, 2, 3$. Therefore $\text{Ext}_{\mathbb{X}}^1(L', \bar{L}(\vec{\omega})) = \text{Hom}_{\mathbb{X}}(\bar{L}, L') = 0$. Since the direct summands of the projective cover of E are $\text{Hom}_{\mathbb{X}}$ -orthogonal, we conclude that $\dim \text{Hom}_{\mathbb{X}}(L', E) = \dim \text{Hom}_{\mathbb{X}}(L', \bar{L}(\vec{\omega})) = 1$ for $L = \bar{L}(\vec{\omega})$, and $\dim \text{Hom}_{\mathbb{X}}(L', \bar{L}(\vec{x})) = 1$ for $L \neq \bar{L}(\vec{\omega})$. ■

As the following example shows, achieving the condition $uv = f\mathbf{1} = vu$ by specialization of a factorization frame for E is not sufficient to obtain a matrix factorization representing E .

EXAMPLE 4.12. Let \mathbb{X} be a weighted projective line of type $(2, 3, 4)$. We consider the almost-split sequence $0 \rightarrow \tau G_2 \rightarrow E_4 \rightarrow G_2 \rightarrow 0$ from Proposition 3.8. This sequence satisfies the assumptions of Lemma 3.5. Hence the factorization frame for E_4 has the shape

$$U_{E_4} = \left[\begin{array}{c|c} \bar{u}_{\tau G_2} & \bar{b} \\ \hline 0 & \bar{u}_{G_2} \end{array} \right], \quad V_{E_4} = \left[\begin{array}{c|c} \bar{v}_{\tau G_2} & \bar{b} \\ \hline 0 & \bar{v}_{G_2} \end{array} \right].$$

If we choose scalars such that $b = 0$, and $u_{\tau G_2}, v_{\tau G_2}$ (resp. u_{G_2}, v_{G_2}) are matrix factorization for τG_2 (respectively G_2), then we obtain a matrix factorization for $\tau G_2 \oplus G_2$, not for E_4 .

In checking the indecomposability of a matrix factorization obtained by specializing matrix frames for bundles of rank two and three, the following observation will be helpful.

OBSERVATION 4.13. *Let \mathbb{X} have a domestic type and E be an indecomposable vector bundle of rank 2 or 3. Then the line bundle summands of $\mathfrak{P}(E)$ are pairwise Hom-orthogonal.*

Proof. For rank two this is a general fact (see Theorem 3.2). For rank three, this follows by inspection of the projective covers for domestic weight triples. ■

5. Matrix factorizations. Throughout this section, we will freely switch from the notation (x_1, x_2, x_3) to (x, y, z) , whenever this is preferable for typographical reasons.

This section presents explicit matrix factorizations for the following cases. Keeping the weight triple (a, b, c) of integers greater than or equal to 2, we first present a general result on the matrix factorizations of the \mathbb{L} -graded triangle singularity $f = x^a + y^b + z^c$ for indecomposable bundles of rank two. Next, we restrict to the weight triple $(2, a, b)$, where we obtain *symmetric* matrix factorizations for indecomposable vector bundles of rank two.

For the second part of the section we restrict to weight triples of domestic case, that is, to the weight triples $(2, 2, n)$, $(2, 3, 3)$, $(2, 3, 4)$ and

(2, 3, 5). Here, we determine *symmetric* matrix factorizations for each indecomposable vector bundle (equivalently, each \mathbb{L} -graded indecomposable Cohen–Macaulay module) of rank ≥ 2 , where for rank two we use the general result for extension bundles. We emphasize that our methods work over any characteristic, yielding matrices whose entries are $\{0, \pm 1\}$ multiples of monomials with small exponents. Still restricting to domestic weight types, we note that the concepts of simple singularities and triangle singularities, as studied in [KST07], [LdIP11], respectively [KLM13], agree exactly for weight type (2, 3, 5). Due to different approaches, the resulting matrix factorizations for $f = x^2 + y^3 + z^5$ are different.

General results. We recall that the group \mathbb{L} acts on $\text{coh}(\mathbb{X})$, and on related mathematical objects like $T = k[x_1, x_2, x_3]$, $S = T/(f)$, $\text{CM}^{\mathbb{L}}(S)$ and $\text{vect}(\mathbb{X})$, by degree shift. The next observation largely simplifies the determination of explicit matrix factorizations.

LEMMA 5.1. *We assume that \mathbb{X} has triple weight type. Let E be a vector bundle admitting a matrix factorization (u, v) . Then for each \vec{x} in \mathbb{L} , also $E(\vec{x})$ admits the matrix factorization (u, v) . In particular, all members of a τ -orbit in $\text{vect}(\mathbb{X})$ admit matrix factorizations by the same matrices.*

Proof. If M is the \mathbb{L} -graded Cohen–Macaulay S -module corresponding to E , and $P_0 \xrightarrow{u} P_1 \xrightarrow{v} P_0 \rightarrow M \rightarrow 0$ is the start of a 2-periodic minimal projective A -resolution for M , then application of the degree shift with \vec{x} yields the start $P_0(\vec{x}) \xrightarrow{u} P_1(\vec{x}) \xrightarrow{v} P_0(\vec{x}) \rightarrow M(\vec{x}) \rightarrow 0$ of a 2-periodic minimal projective A -resolution for $M(\vec{x})$, just keeping the matrices u and v . ■

Assume that \mathbb{X} is a weighted projective line of triple weight type (a, b, c) , not necessarily domestic. We recall from [KLM13, Theorem 4.2] that each indecomposable vector bundle E of rank two is an extension bundle, that is, it is the middle term $E_L\langle\vec{x}\rangle$ of ‘the’ non-split exact sequence $0 \rightarrow L(\vec{\omega}) \rightarrow E \rightarrow L(\vec{x}) \rightarrow 0$ for some line bundle L and some $\vec{0} \leq \vec{x} \leq \vec{\delta}$, where $\vec{\delta} = \vec{c} + 2\vec{\omega}$. By Lemma 5.1 we obtain matrix factorizations $(u_{\vec{x}}, v_{\vec{x}})$ for $E_L\langle\vec{x}\rangle$ where the matrices $u_{\vec{x}}$ and $v_{\vec{x}}$ are independent of L . To see this, one uses formula (3.2). In the following, we are going to construct matrix factorizations for many vector bundles E (or Cohen–Macaulay modules M). In order to describe such a matrix factorization uniquely, we list the projective cover $\mathfrak{P}(E)$ of E (or M) together with the matrix pair (u, v) of the factorization. If we represent $\mathfrak{P}(E)$ as a direct sum of line bundles $\mathcal{O}(\vec{y}_j)$, $j = 1, \dots, s$, then the triple notation $(u, v, \mathfrak{P}(E))$, or equivalently $(u, v, (\vec{y}_j))$, determines E (or M) uniquely, up to isomorphism.

PROPOSITION 5.2. *Let \mathbb{X} be a weighted projective line of triple weight type (a, b, c) . Then the extension bundle $E = E_L\langle\vec{x}\rangle$, where $\vec{x} = \sum_{i=1}^3 l_i \vec{x}_i$ and $\vec{0} \leq \vec{x} \leq \vec{\delta}$, admits the matrix factorization $(u_{\vec{x}}, v_{\vec{x}}, (\vec{\omega}, \vec{x} - (1 + l_i \vec{x}_i)_{i=1,2,3}))$,*

where

$$u_{\vec{x}} = \begin{bmatrix} 0 & x^{(1+l_1)} & y^{(1+l_2)} & z^{(1+l_3)} \\ x^{(1+l_1)} & 0 & z^{c-(1+l_3)} & -y^{b-(1+l_2)} \\ y^{(1+l_2)} & -z^{c-(1+l_3)} & 0 & x^{a-(1+l_1)} \\ z^{(1+l_3)} & y^{b-(1+l_2)} & -x^{a-(1+l_1)} & 0 \end{bmatrix},$$

$$v_{\vec{x}} = \begin{bmatrix} 0 & x^{a-(1+l_1)} & y^{b-(1+l_2)} & z^{c-(1+l_3)} \\ x^{a-(1+l_1)} & 0 & -z^{(1+l_3)} & y^{(1+l_2)} \\ y^{b-(1+l_2)} & z^{(1+l_3)} & 0 & -x^{(1+l_1)} \\ z^{c-(1+l_3)} & -y^{(1+l_2)} & x^{(1+l_1)} & 0 \end{bmatrix}.$$

Proof. From Theorem 3.2, the projective cover of the extension bundle $E_L\langle\vec{x}\rangle$ is given by $\mathfrak{P}(E_L\langle\vec{x}\rangle) = L(\vec{\omega}) \oplus \bigoplus_{i=1}^3 L(\vec{x} - (1+l_i)\vec{x}_i)$.

The corresponding factorization frame for $E_L\langle\vec{x}\rangle$ has the shape

$$U_{\vec{x}} = \begin{bmatrix} 0 & x^{(1+l_1)} & y^{(1+l_2)} & z^{(1+l_3)} \\ x^{(1+l_1)} & 0 & z^{c-(1+l_3)} & y^{b-(1+l_2)} \\ y^{(1+l_2)} & z^{c-(1+l_3)} & 0 & x^{a-(1+l_1)} \\ z^{(1+l_3)} & y^{b-(1+l_2)} & x^{a-(1+l_1)} & 0 \end{bmatrix},$$

$$V_{\vec{x}} = \begin{bmatrix} 0 & x^{a-(1+l_1)} & y^{b-(1+l_2)} & z^{c-(1+l_3)} \\ x^{a-(1+l_1)} & 0 & z^{(1+l_3)} & y^{(1+l_2)} \\ y^{b-(1+l_2)} & z^{(1+l_3)} & 0 & x^{(1+l_1)} \\ z^{c-(1+l_3)} & y^{(1+l_2)} & x^{(1+l_1)} & 0 \end{bmatrix}.$$

From Lemmas 4.10 and 4.11 we need to choose non-zero scalars such that $u_{\vec{x}}v_{\vec{x}} = f\mathbb{1} = v_{\vec{x}}u_{\vec{x}}$ and $(u_{\vec{x}}, v_{\vec{x}})$ is indecomposable. By Theorem 3.2 the line bundle summands of the projective cover of $E_L\langle\vec{x}\rangle$ are Hom-orthogonal; it is then easy to check that the above choice of scalars yields an indecomposable matrix factorization. That we get, indeed, a matrix factorization attached to E then follows from Proposition 3.11. ■

We next assume weight type $(2, a, b)$ and show that each extension bundle $E = E_L\langle\vec{x}\rangle$ admits a *symmetric* matrix factorization. We recall from Theorem 3.3 that the suspension functor [1] for $\text{vect}(\mathbb{X})$ is induced by the degree shift $E \mapsto E(\vec{x}_1)$. The minimal projective resolution of E has the form

$$(5.1) \quad \mathfrak{P}(E)(-\vec{c}) \xrightarrow{\bar{v}=\bar{u}(-\vec{x}_1)} \mathfrak{P}(E)(-\vec{x}_1) \xrightarrow{\bar{u}} \mathfrak{P}(E) \rightarrow E \rightarrow 0.$$

Interpreting the above sequence on the level of S -modules shows that the symmetry condition $\bar{v} = \bar{u}(-\vec{x}_1)$ is satisfied over $S = T/(f)$. It remains to lift the maps \bar{u} and \bar{v} to T -linear maps u and v such that $v = u(-\vec{x}_1)$, and moreover the matrix factorization identity $u(-\vec{x}_1)u = f\mathbf{1}$ holds.

PROPOSITION 5.3. *Let \mathbb{X} be a weighted projective line of triple weight type $(2, a, b)$. Then each extension bundle $E = E_L\langle\vec{x}\rangle$, where $\vec{x} = l_2\vec{x}_2 + l_3\vec{x}_3$ and $0 \leq l_2 \leq a - 2$, $0 \leq l_3 \leq b - 2$, admits a symmetric matrix factorization $(u_{\vec{x}}, v_{\vec{x}}, (\vec{\omega}, \vec{x} - \vec{x}_1, l_3\vec{x}_3 - \vec{x}_2, l_2\vec{x}_2 - \vec{x}_3))$ of $f = x^2 + y^a + z^b$, where*

$$u_{\vec{x}} = v_{\vec{x}} = \begin{bmatrix} x & 0 & -z^{b-l_3-1} & y^{a-l_2-1} \\ 0 & x & y^{l_2+1} & z^{l_3+1} \\ -z^{l_3+1} & y^{a-l_2-1} & -x & 0 \\ y^{l_2+1} & z^{b-l_3-1} & 0 & -x \end{bmatrix}.$$

Proof. We fix a decomposition of $\mathfrak{P}(E)$ into line bundles that is transferred to the other terms by degree shifts with $-\vec{x}_1$ and $-\vec{c}$. We then obtain a matrix factorization $(u_{\vec{x}}, v_{\vec{x}})$ such that $u_{\vec{x}}$ and $v_{\vec{x}} = u_{\vec{x}}(-\vec{x}_1)$ are represented by the same matrix. Defining the above specialization $u_{\vec{x}}$ of the factorization frame for E , we achieve the matrix factorization condition $u_{\vec{x}}^2 = f\mathbf{1}$ over T . Using arguments as before, it is again easy to check that this matrix factorization has a trivial endomorphism algebra. Hence it is indecomposable, and so by Proposition 3.12 it represents E . ■

The domestic case. The main result of this section, and actually the main result of this paper, concerns the domestic case, necessarily of type $(2, a, b)$, where, for each indecomposable vector bundle of rank at least two, we determine explicitly a *symmetric* matrix factorization.

THEOREM 5.4. *We assume a triangle singularity $f = x^2 + y^a + z^b$ of domestic type. For each indecomposable bundle E of rank at least two, we obtain a symmetric matrix factorization $u^2 = f\mathbf{1}$ of f representing E . The matrix entries of u are scalar multiples of monomials in x, y, z with small exponents. Moreover, the scalars may be taken from $\{0, \pm 1\}$.*

Proof. Interpreting the minimal projective resolution (5.1) as a sequence of \mathbb{L} -graded Cohen–Macaulay modules over S , we lift it to a matrix factorization of f over T , thus obtaining a commutative diagram

$$(5.2) \quad \begin{array}{ccccccccc} \cdots & \longrightarrow & P_0(-\vec{c}) & \xrightarrow{v} & P_0(-\vec{x}_1) & \xrightarrow{u} & P_0 & \xrightarrow{\pi} & M & \longrightarrow & 0 \\ & & \nu \downarrow & & \nu \downarrow & & \nu \downarrow & & \parallel & & \\ \cdots & \longrightarrow & \bar{P}_0(-\vec{c}) & \xrightarrow{\bar{v}=\bar{u}(-\vec{x}_1)} & \bar{P}_0(-\vec{x}_1) & \xrightarrow{\bar{u}} & \bar{P}_0 & \xrightarrow{\bar{\pi}} & M & \longrightarrow & 0 \end{array}$$

First we are going to show that the matrix factorization (u, v) is symmetric, that is, $v = u(-\vec{x}_1)$. Now, reduction modulo (f) sends v and $u(-\vec{x}_1)$ to

the same map \bar{v} . Since \mathbb{X} has domestic type, and by invoking compatible decompositions of the P_i and \bar{P}_i into indecomposable projectives, Theorem 4.3 shows that $v = u(-\bar{x}_1)$. Moreover, by the same theorem, the matrix entries (with respect to the chosen decomposition) are scalar multiples of monomials with small exponents.

That the scalars, actually, can be chosen from among 0 and ± 1 follows through a case-by-case analysis from the following propositions. ■

For the rest of this section, we derive explicit matrix factorizations for all indecomposable vector bundles of rank at least two. We recall (see Lemma 5.1) that a single matrix factorization is sufficient to represent all members of a fixed Auslander–Reiten orbit.

The triangle singularity $x^2 + y^2 + z^n$ ($n \geq 2$). Here, the projective covers are given by Proposition 3.6. Since for type $(2, 2, n)$ all indecomposable vector bundles that are not line bundles have rank two, the corresponding matrix factorizations are a special case of Proposition 5.3.

PROPOSITION 5.5. *Let E_i denote the extension bundle $E_L\langle i\bar{x}_3 \rangle$ for $i = 0, \dots, n-2$. Then the bundle E_i yields a symmetric matrix factorization $(u_{E_i}, v_{E_i}, \mathfrak{P}(E_i))$ of $x^2 + y^2 + z^n$ as follows:*

$$u_{E_i} = v_{E_i} = \begin{bmatrix} x & 0 & -z^{n-i-1} & y \\ 0 & x & y & z^{i+1} \\ -z^{i+1} & y & -x & 0 \\ y & z^{n-i-1} & 0 & -x \end{bmatrix}$$

for $i = 0, \dots, n-2$. ■

The triangle singularity $x^2 + y^3 + z^3$. We will use the projective covers described in Proposition 3.7 and use the notation from the Auslander–Reiten quiver of type $(2, 3, 3)$ depicted there.

PROPOSITION 5.6. *For the singularity $x^2 + y^3 + z^3$ we obtain the symmetric matrix factorizations $(u_2, v_2, \mathfrak{P}(E_2))$, $(u_2, v_2, \mathfrak{P}(F_2))$, $(u_2, v_2, \mathfrak{P}(G_2))$, $(u_3, v_3, \mathfrak{P}(E_3))$, where*

$$u_2 = v_2 = \begin{bmatrix} x & 0 & -z^2 & y^2 \\ 0 & x & y & z \\ -z & y^2 & -x & 0 \\ y & z^2 & 0 & -x \end{bmatrix}, \quad u_3 = v_3 = \begin{bmatrix} x & yz & 0 & y^2 & 0 & -z^2 \\ 0 & -x & z & 0 & -y & 0 \\ 0 & z^2 & x & yz & 0 & y^2 \\ y & 0 & 0 & -x & z & 0 \\ 0 & -y^2 & 0 & z^2 & x & yz \\ -z & 0 & y & 0 & 0 & -x \end{bmatrix}.$$

Proof. The vector bundles E_2, F_2, G_2 are extension bundles determined by the pairs $(\mathcal{O}(-\vec{\omega}), \vec{0})$, $(\mathcal{O}(\vec{x}_2 + \vec{\omega}), \vec{0})$, $(\mathcal{O}(\vec{x}_3 + \vec{\omega}), \vec{0})$, respectively. Therefore the claim for those bundles results from Proposition 5.3. Concerning the vector bundle E_3 , it is first checked that $u_3 v_3 = f\mathbf{1} = v_3 u_3$. Since, moreover, the direct summands of $\mathfrak{P}(E_3)$ are mutually Hom-orthogonal, it is easy to check that the matrix factorization (u_3, v_3) is indecomposable. By Proposition 3.11 it then represents the vector bundle in question. ■

The triangle singularity $x^2 + y^3 + z^4$. For weight type $(2, 3, 4)$, each indecomposable vector bundle is of rank 1, 2, 3 or 4. As in Proposition 5.3, and using the notation introduced there, we only need to determine matrix factorizations for the vector bundles E_2, E_3, E_4 and F_2 , since by symmetry E_i and $E_i(\vec{x}_1 - 2\vec{x}_3)$ will yield the same matrix pair. Moreover, matrix factorizations for the rank-two bundles E_2 and G_2 are already given by Proposition 5.3. We thus obtain:

PROPOSITION 5.7. *For the singularity $x^2 + y^3 + z^4$ we obtain the symmetric matrix factorizations $(u_{E_i}, v_{E_i}, \mathfrak{P}(E_i))$ for $i = 2, 3, 4$, $(u_{E_i}, v_{E_i}, \mathfrak{P}(E_i(\vec{x}_1 - 2\vec{x}_3)))$ for $i = 2, 3$, $(u_{G_2}, v_{G_2}, \mathfrak{P}(G_2))$, where*

$$u_{E_2} = v_{E_2} = \begin{bmatrix} x & 0 & -z^3 & y^2 \\ 0 & x & y & z \\ -z & y^2 & -x & 0 \\ y & z^3 & 0 & -x \end{bmatrix},$$

$$u_{E_3} = v_{E_3} = \begin{bmatrix} x & 0 & z^3 & 0 & -y^2 & -yz^2 \\ 0 & x & yz & 0 & z^2 & -y^2 \\ z & 0 & -x & y & 0 & 0 \\ 0 & 0 & y^2 & x & yz & z^3 \\ -y & z^2 & 0 & 0 & -x & 0 \\ 0 & -y & 0 & z & 0 & -x \end{bmatrix},$$

$$u_{E_4} = v_{E_4} = \left[\begin{array}{cccc|cccc} x & 0 & -z^2 & y^2 & 0 & -yz & 0 & 0 \\ 0 & x & y & z^2 & z & 0 & 0 & 0 \\ -z^2 & y^2 & -x & 0 & 0 & 0 & 0 & -yz \\ y & z^2 & 0 & -x & 0 & 0 & z & 0 \\ \hline 0 & 0 & 0 & 0 & -x & 0 & -z^2 & y^2 \\ 0 & 0 & 0 & 0 & 0 & -x & y & z^2 \\ 0 & 0 & 0 & 0 & -z^2 & y^2 & x & 0 \\ 0 & 0 & 0 & 0 & y & z^2 & 0 & x \end{array} \right],$$

$$u_{G_2} = v_{G_2} = \begin{bmatrix} x & 0 & -z & y^2 \\ 0 & x & y & z^2 \\ -z^2 & y^2 & -x & 0 \\ y & z^2 & 0 & -x \end{bmatrix}.$$

Proof. It is easy to check that the above matrices satisfy the matrix equation $u^2 = f\mathbb{1}$. By Proposition 3.11, it remains to prove the indecomposability of those matrix factorizations by showing that their endomorphism rings are trivial. For the indecomposable vector bundles of rank two and three, the indecomposability follows easily, since the indecomposable direct summands of their projective covers are Hom-orthogonal. It remains to check indecomposability for (u_{E_4}, v_{E_4}) , where

$$u_{E_4} = v_{E_4} = \left[\begin{array}{c|c} u_{\tau_X T} & B \\ \hline 0 & u_T \end{array} \right],$$

by using the explicit form of the projective cover of E_4 by means of Proposition 3.8. The same argument yields the shape of an endomorphism (K, H) for (u_{E_4}, v_{E_4}) :

$$H = \left[\begin{array}{c|c} H_1 & 0 \\ \hline H_3 & H_4 \end{array} \right] = \begin{bmatrix} f_1 & 0 & 0 & 0 & | & 0 & 0 & 0 & 0 \\ 0 & f_2 & 0 & 0 & | & 0 & 0 & 0 & 0 \\ 0 & 0 & f_3 & 0 & | & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & f_4 & | & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & f_{5,4}z & | & f_5 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & | & 0 & f_6 & 0 & 0 \\ 0 & f_{7,2}z & 0 & 0 & | & 0 & 0 & f_7 & 0 \\ 0 & 0 & 0 & 0 & | & 0 & 0 & 0 & f_8 \end{bmatrix},$$

$$K = \left[\begin{array}{c|c} K_1 & 0 \\ \hline K_3 & K_4 \end{array} \right] = \begin{bmatrix} g_1 & 0 & 0 & 0 & | & 0 & 0 & 0 & 0 \\ 0 & g_2 & 0 & 0 & | & 0 & 0 & 0 & 0 \\ 0 & 0 & g_3 & 0 & | & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & g_4 & | & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & g_{5,4}z & | & g_5 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & | & 0 & g_6 & 0 & 0 \\ 0 & g_{7,2}z & 0 & 0 & | & 0 & 0 & g_7 & 0 \\ 0 & 0 & 0 & 0 & | & 0 & 0 & 0 & g_8 \end{bmatrix}.$$

It follows that $Hu_{E_4} = u_{E_4}K$. Now, in block matrix form we have

$$\begin{aligned} 0 &= \left[\begin{array}{c|c} H_1 & 0 \\ \hline H_3 & H_4 \end{array} \right] \left[\begin{array}{c|c} u_{\tau_{\mathbb{X}}T} & B \\ \hline 0 & u_T \end{array} \right] - \left[\begin{array}{c|c} u_{\tau_{\mathbb{X}}T} & B \\ \hline 0 & u_T \end{array} \right] \left[\begin{array}{c|c} K_1 & 0 \\ \hline K_3 & K_4 \end{array} \right] \\ &= \left[\begin{array}{c|c} H_1u_{\tau_{\mathbb{X}}T} - u_{\tau_{\mathbb{X}}T}K_1 & H_1B - BK_4 \\ \hline H_3u_{\tau_{\mathbb{X}}T} - u_TK_3 & H_3B + H_4u_T - u_TK_4 \end{array} \right], \end{aligned}$$

and we deduce that (K_1, H_1) is an endomorphism for $(u_{\tau_{\mathbb{X}}T}, v_{\tau_{\mathbb{X}}T})$. Therefore

$$H_1 = K_1 = \lambda \mathbf{1}_4.$$

Moreover

$$\begin{aligned} 0 &= H_3u_{\tau_{\mathbb{X}}T} - u_TK_3 \\ &= \begin{bmatrix} yzf_{5,4} & z^3f_{5,4} + z^3g_{7,2} & 0 & -xzf_{5,4} + xzg_{5,4} \\ 0 & -yzg_{7,2} & 0 & 0 \\ 0 & xzf_{7,2} - xzg_{7,2} & yzf_{7,2} & z^3f_{7,2} + z^3g_{5,4} \\ 0 & 0 & 0 & -yzg_{5,4} \end{bmatrix}. \end{aligned}$$

Thus $K_3 = H_3 = 0$, hence (K_4, H_4) is an endomorphism for (u_T, v_T) , and from the indecomposability of (u_T, v_T) we get

$$H_4 = K_4 = \mu \mathbf{1}_4.$$

The equation $H_1B = BK_4$ implies that $\lambda = \mu$. Therefore (u_{E_4}, v_{E_4}) is indecomposable, and the claim follows from Proposition 3.11. ■

The triangle singularity $x^2 + y^3 + z^5$. In this case, each indecomposable vector bundle on \mathbb{X} is of rank m , $1 \leq m \leq 6$, and there is a single $\tau_{\mathbb{X}}$ -orbit of vector bundles of rank 5, and a single one of rank 6.

For a representative system of indecomposable vector bundles $E_2, E_3, \dots, E_6, F_2, F_4$ and G_3 of rank at least two, we use the choices and notation of Section 3.

PROPOSITION 5.8. *For the singularity $f = x^2 + y^3 + z^5$ we obtain eight matrix factorizations: $(u_{E_2}, v_{E_2}, \mathfrak{P}(E_2)), \dots, (u_{E_6}, v_{E_6}, \mathfrak{P}(E_6))$, and $(u_{F_2}, v_{F_2}, \mathfrak{P}(F_2)), (u_{F_4}, v_{F_4}, \mathfrak{P}(F_4))$, and $(u_{G_3}, v_{G_3}, \mathfrak{P}(G_3))$, where*

$$u_{E_2} = v_{E_2} = \begin{bmatrix} x & 0 & -z^4 & y^2 \\ 0 & x & y & z \\ -z & y^2 & -x & 0 \\ y & z^4 & 0 & -x \end{bmatrix},$$

$$u_{E_3} = v_{E_3} = \begin{bmatrix} x & 0 & -y^2 & 0 & z^4 & -yz^3 \\ 0 & x & yz & 0 & y^2 & z^4 \\ -y & 0 & -x & z^3 & 0 & 0 \\ 0 & 0 & z^2 & x & yz & -y^2 \\ z & y & 0 & 0 & -x & 0 \\ 0 & z & 0 & -y & 0 & -x \end{bmatrix},$$

$$u_{E_4} = v_{E_4} = \left[\begin{array}{cccc|cccc} x & 0 & z^3 & y^2 & 0 & 0 & 0 & yz \\ 0 & x & -y & z^2 & 0 & 0 & z & 0 \\ z^2 & -y^2 & -x & 0 & 0 & yz & 0 & 0 \\ y & z^3 & 0 & -x & z & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & x & 0 & -z^3 & -y^2 \\ 0 & 0 & 0 & 0 & 0 & x & y & -z^2 \\ 0 & 0 & 0 & 0 & -z^2 & y^2 & -x & 0 \\ 0 & 0 & 0 & 0 & -y & -z^3 & 0 & -x \end{array} \right],$$

$$u_{E_5} = v_{E_5} = \left[\begin{array}{cccccc|cccc} x & 0 & 0 & y^2 & yz^2 & z^4 & 0 & 0 & 0 & -z^3 \\ 0 & x & 0 & -z^3 & y^2 & yz^2 & 0 & 0 & 0 & -yz \\ 0 & 0 & x & -yz & -z^3 & y^2 & 0 & 0 & z^2 & 0 \\ y & -z^2 & 0 & -x & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & y & -z^2 & 0 & -x & 0 & z & 0 & 0 & 0 \\ z & 0 & y & 0 & 0 & -x & 0 & z^2 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & x & 0 & z^3 & y^2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & x & -y & z^2 \\ 0 & 0 & 0 & 0 & 0 & 0 & z^2 & -y^2 & -x & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & y & z^3 & 0 & -x \end{array} \right],$$

$$u_{E_6} = v_{E_6} = \left[\begin{array}{cccccc|cccccc} x & 0 & 0 & y^2 & yz^2 & z^4 & 0 & -yz & 0 & 0 & 0 & 0 \\ 0 & x & 0 & -z^3 & y^2 & yz^2 & z^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x & -yz & -z^3 & y^2 & y & 0 & 0 & 0 & 0 & 0 \\ y & -z^2 & 0 & -x & 0 & 0 & 0 & 0 & 0 & 0 & yz & z^3 \\ 0 & y & -z^2 & 0 & -x & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ z & 0 & y & 0 & 0 & -x & 0 & 0 & 0 & -y & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & -x & 0 & 0 & y^2 & yz^2 & z^4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -x & 0 & -z^3 & y^2 & yz^2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -x & -yz & -z^3 & y^2 \\ 0 & 0 & 0 & 0 & 0 & 0 & y & -z^2 & 0 & x & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & y & -z^2 & 0 & x & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & z & 0 & y & 0 & 0 & x \end{array} \right],$$

$$\begin{aligned}
u_{F_2} = v_{F_2} &= \begin{bmatrix} x & 0 & -z^3 & y^2 \\ 0 & x & y & z^2 \\ -z^2 & y^2 & -x & 0 \\ y & z^3 & 0 & -x \end{bmatrix}, \\
u_{F_4} = v_{F_4} &= \left[\begin{array}{cccc|cccc} x & 0 & z^3 & y^2 & 0 & yz^2 & 0 & 0 \\ 0 & x & -y & z^2 & z & 0 & 0 & 0 \\ z^2 & -y^2 & -x & 0 & 0 & 0 & 0 & yz \\ y & z^3 & 0 & -x & 0 & 0 & z^2 & 0 \\ \hline 0 & 0 & 0 & 0 & -x & 0 & -z^3 & y^2 \\ 0 & 0 & 0 & 0 & 0 & -x & -y & -z^2 \\ 0 & 0 & 0 & 0 & -z^2 & -y^2 & x & 0 \\ 0 & 0 & 0 & 0 & y & -z^3 & 0 & x \end{array} \right], \\
u_{G_3} = v_{G_3} &= \begin{bmatrix} x & 0 & 0 & y^2 & yz^2 & z^4 \\ 0 & x & 0 & -z^3 & y^2 & yz^2 \\ 0 & 0 & x & -yz & -z^3 & y^2 \\ y & -z^2 & 0 & -x & 0 & 0 \\ 0 & y & -z^2 & 0 & -x & 0 \\ z & 0 & y & 0 & 0 & -x \end{bmatrix}.
\end{aligned}$$

Proof. Similarly to the case $(2, 3, 4)$ we check that $uv = f\mathbf{1} = vu$ and verify that these matrix factorizations are indecomposable. The case of rank-two bundles is covered by Proposition 5.3. The matrix factorizations for the rank-three bundles E_3 and F_3 are verified following the arguments of Proposition 5.7. Concerning the remaining vector bundles, we determine the pairs (u_{E_i}, v_{E_i}) for $i = 4, 5, 6$ and (u_{F_i}, v_{F_i}) for $i = 4$ by specialization of factorization frames coming from distinguished exact sequences by means of Proposition 3.9. In particular, for (u_{E_4}, v_{E_4}) , (u_{E_5}, v_{E_5}) , (u_{E_6}, v_{E_6}) and (u_{F_4}, v_{F_4}) , we use the exact sequences $0 \rightarrow \tau_{\mathbb{X}}^2 F_2 \rightarrow E_4 \rightarrow \tau_{\mathbb{X}}^{-2} F_2 \rightarrow 0$, $0 \rightarrow \tau_{\mathbb{X}} G_3 \rightarrow E_5 \rightarrow \tau_{\mathbb{X}}^- F_2 \rightarrow 0$, $0 \rightarrow G_3 \rightarrow E_6 \rightarrow \tau_{\mathbb{X}}^- G_3 \rightarrow 0$ and $0 \rightarrow \tau_{\mathbb{X}} F_2 \rightarrow F_4 \rightarrow F_2 \rightarrow 0$. ■

REMARK 5.9. Observe that $0 \rightarrow \tau_{\mathbb{X}} F_4 \rightarrow E_6 \rightarrow \tau_{\mathbb{X}}^- F_2 \rightarrow 0$ is a distinguished exact sequence, thus satisfying the assumptions of Lemma 3.5. Using the resulting direct decomposition $\mathfrak{P}(E_6) = \mathfrak{P}(\tau F_4) \oplus \mathfrak{P}(\tau^- F_2)$, we get another—essentially different—pair of matrices, also yielding a matrix factorization of E_6 :

$$u'_{E_6} = v'_{E_6} = \left[\begin{array}{cccccccc|cccc} x & 0 & z^3 & y^2 & 0 & yz^2 & 0 & 0 & 0 & 0 & 0 & -z^3 \\ 0 & x & -y & z^2 & z & 0 & 0 & 0 & 0 & 0 & 0 & y \\ z^2 & -y^2 & -x & 0 & 0 & 0 & 0 & yz & y & z^3 & 0 & 0 \\ y & z^3 & 0 & -x & 0 & 0 & z^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x & 0 & -z^3 & y^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -x & -y & -z^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -z^2 & -y^2 & x & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & y & -z^3 & 0 & x & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x & 0 & z^3 & y^2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x & -y & z^2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & z^2 & -y^2 & -x & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & y & z^3 & 0 & -x \end{array} \right].$$

In particular, the numbers of zero entries differ for both matrix factorizations.

REMARK 5.10. If we compare our matrix factorizations for E_6 with the matrix factorization obtained in [KST07], we also see that they are essentially different, since for one matrix factorization there appear monomial entries z^4 , for the other one there do not.

6. Appendix: Tables of projective covers. The figures of this section yield compact visual information on the projective covers of indecomposable vector bundles of rank at least two. Our figures may be especially useful for specialists from the representation theory of finite-dimensional algebras investigating the related situation in preprojective, or preinjective, components for tame concealed quivers. We note that, in the representation-theoretic context, the line bundle notation $\mathcal{O}(\vec{x})$, reduced in the figures to (\vec{x}) , is not established, so the given positions in the mesh category of the associated extended Dynkin quiver should be useful. The names E_i , F_j and G_l for selected vector bundles are those from Section 3.

Weight type $(2, 2, n)$.




Fig. 1. $(2, 2, n)$: Projective cover of $E = E(i\vec{x}_3)$, where $\vec{u} = \vec{x}_1 - \vec{x}_2$

Weight type (2, 3, 3). The projective covers of the ‘remaining’ indecomposable vector bundles are obtained by applying twice the rotation $X \mapsto X(\vec{x}_2 - \vec{x}_3)$ around the central axis.




Fig. 2. (2, 3, 3): Projective cover of F_2 and E_3

Weight type (2, 3, 4). The next figure yields the projective covers for the indecomposable vector bundles E_2 , E_3 , E_4 and G_2 . For the projective cover of E_4 one has to combine the projective covers of G_2 and τG_2 . By means of reflection in the central horizontal axis $X \mapsto X(\vec{x}_1 - 2\vec{x}_3)$ one obtains the projective covers for the ‘missing’ vector bundles $E_2(\vec{x}_1 - 2\vec{x}_3)$ and $E_3(\vec{x}_1 - 2\vec{x}_3)$.




Fig. 3. (2, 3, 4): Projective covers for E_2 , E_3 , E_4 , G_2 and τG_2

Here and below, by \star (resp. \blacklozenge) we have marked the line bundle summands of the projective covers of τG_2 (resp. G_2); together they form the projective cover of E_4 .

Weight type (2, 3, 5).




Fig. 4 (2, 3, 5): Projective covers for $E_2, E_3, E_4, E_5, E_6, F_2, F_4$ and G_3 (continued next page)




Fig. 4 (cont.)

Acknowledgements. The first author was supported by the Research Project from the MNiSW specific subsidy for young scientists of the Department of Mathematics and Physics of Szczecin University 504-4000-240-835.

The third author was supported by the Polish Scientific Grant Narodowe Centrum Nauki DEC-2011/01/B/ST1/06469.

The authors further want to thank the referee for helpful suggestions and a critical checking of the manuscript.

REFERENCES

- [Buc86] R.-O. Buchweitz, *Maximal Cohen–Macaulay modules and Tate-cohomology over Gorenstein rings*, preprint, 1986; <http://hdl.handle.net/1807/16682>.
- [Eis80] D. Eisenbud, *Homological algebra on a complete intersection, with an application to group representations*, Trans. Amer. Math. Soc. 260 (1980), 35–64.
- [GL87] W. Geigle and H. Lenzing, *A class of weighted projective curves arising in representation theory of finite-dimensional algebras*, in: Singularities, Representation of Algebras, and Vector Bundles (Lambrech, 1985), Lecture Notes in Math. 1273, Springer, Berlin, 1987, 265–297.
- [GL91] W. Geigle and H. Lenzing, *Perpendicular categories with applications to representations and sheaves*, J. Algebra 144 (1991), 273–343.
- [Hap88] D. Happel, *Triangulated Categories in the Representation Theory of Finite-Dimensional Algebras*, London Math. Soc. Lecture Note Ser. 119, Cambridge Univ. Press, Cambridge, 1988.
- [Hüb96] T. Hübner, *Exzeptionelle Vektorbündel und Reflektionen an Kippgarben über projektiven gewichteten Kurven*, Dissertation, Univ. Paderborn, 1996.
- [KST07] H. Kajiura, K. Saito, and A. Takahashi, *Matrix factorization and representations of quivers. II. Type ADE case*, Adv. Math. 211 (2007), 327–362.
- [KLM13] D. Kussin, H. Lenzing, and H. Meltzer, *Triangle singularities, ADE-chains, and weighted projective lines*, Adv. Math. 237 (2013), 194–251.
- [LdlP11] H. Lenzing and J. A. de la Peña, *Extended canonical algebras and Fuchsian singularities*, Math. Z. 268 (2011), 143–167.

- [LM92] H. Lenzing and H. Meltzer, *Sheaves on a weighted projective line of genus one, and representations of a tubular algebra*, in: Representations of Algebras (Ottawa, ON, 1992), CMS Conf. Proc. 14, Amer. Math. Soc., Providence, RI, 1993, 313–337.
- [LR] H. Lenzing and S. Q. Ruan, *Vector bundles of rank two on a weighted projective line*, in preparation.
- [Mel04] H. Meltzer, *Exceptional vector bundles, tilting sheaves and tilting complexes for weighted projective lines*, Mem. Amer. Math. Soc. 171 (2004), no. 808.
- [Orl09] D. Orlov, *Derived categories of coherent sheaves and triangulated categories of singularities*, in: Algebra, Arithmetic, and Geometry: In Honor of Yu. I. Manin. Vol. II, Progr. Math. 270, Birkhäuser Boston, Boston, MA, 2009, 503–531.
- [Yos90] Y. Yoshino, *Cohen–Macaulay Modules over Cohen–Macaulay Rings*, London Math. Soc. Lecture Note Ser. 146, Cambridge Univ. Press, Cambridge, 1990.

Dawid Edmund Kędzierski, Hagen Meltzer
Instytut Matematyki
Uniwersytet Szczeciński
70-451 Szczecin, Poland
E-mail: d.e.kedzierski@wp.pl
meltzer@wmf.univ.szczecin.pl

Helmut Lenzing
Institut für Mathematik
Universität Paderborn
33095 Paderborn, Germany
E-mail: helmut@math.uni-paderborn.de

*Received 19 November 2013;
revised 26 January 2015*

(6076)